



Eastern Whip-poor-will

(Antrostomus vociferous) in Ontario

Ontario Recovery Strategy Series

Draft

2019

About the Ontario Recovery Strategy Series

This series presents the collection of recovery strategies that are prepared or adopted as advice to the Province of Ontario on the recommended approach to recover species at risk. The Province ensures the preparation of recovery strategies to meet its commitments to recover species at risk under the *Endangered Species Act 2007* (ESA) and the Accord for the Protection of Species at Risk in Canada.

What is recovery?

Recovery of species at risk is the process by which the decline of an endangered, threatened, or extirpated species is arrested or reversed, and threats are removed or reduced to improve the likelihood of a species' persistence in the wild.

What is a recovery strategy?

Under the ESA a recovery strategy provides the best available scientific knowledge on what is required to achieve recovery of a species. A recovery strategy outlines the habitat needs and the threats to the survival and recovery of the species. It also makes recommendations on the objectives for protection and recovery, the approaches to achieve those objectives, and the area that should be considered in the development of a habitat regulation. Sections 11 to 15 of the ESA outline the required content and timelines for developing recovery strategies published in this series.

Recovery strategies are required to be prepared for endangered and threatened species within one or two years respectively of the species being added to the Species at Risk in Ontario list. Recovery strategies are required to be prepared for extirpated species only if reintroduction is considered feasible.

What's next?

Nine months after the completion of a recovery strategy a government response statement will be published which summarizes the actions that the Government of Ontario intends to take in response to the strategy. The implementation of recovery strategies depends on the continued cooperation and actions of government agencies, individuals, communities, land users, and conservationists.

For more information

To learn more about species at risk recovery in Ontario, please visit the Ministry of Environment, Conservation and Parks Species at Risk webpage at: www.ontario.ca/speciesatrisk

1 **Recommended citation**

2 Ministry of the Environment, Conservation and Parks. 2019. DRAFT Recovery Strategy
3 for the Eastern Whip-poor-will (*Antrostomus vociferus*) in Ontario. Ontario Recovery
4 Strategy Series. Prepared by the Ministry of the Environment, Conservation and Parks,
5 Peterborough, Ontario. iv + 6 pp. + Appendix. Adoption of the Recovery Strategy for
6 Eastern Whip-poor-will (*Antrostomus vociferus*) in Canada (Environment and Climate
7 Change Canada 2018).

8 © Queen's Printer for Ontario, 2019
9 ISBN [*MECP will insert prior to final publication.*]

10 Content (excluding illustrations) may be used without permission with appropriate credit
11 to the source, except where use of an image or other item is prohibited in the content
12 use statement of the adopted federal recovery strategy.

13 Cette publication hautement spécialisée « Recovery strategies prepared under the
14 *Endangered Species Act, 2007* », n'est disponible qu'en anglais en vertu du Règlement
15 411/97 qui en exempte l'application de la [Loi sur les services en français](#). Pour obtenir
16 de l'aide en français, veuillez communiquer avec recovery.planning@ontario.ca.

17

18 **Declaration**

19 The recovery strategy for the Eastern Whip-poor-will (*Antrostomus vociferus*) was
20 developed in accordance with the requirements of the *Endangered Species Act, 2007*
21 (ESA). This recovery strategy has been prepared as advice to the Government of
22 Ontario, other responsible jurisdictions and the many different constituencies that may
23 be involved in recovering the species.

24 The recovery strategy does not necessarily represent the views of all individuals who
25 provided advice or contributed to its preparation, or the official positions of the
26 organizations with which the individuals are associated.

27 The recommended goals, objectives and recovery approaches identified in the strategy
28 are based on the best available knowledge and are subject to revision as new
29 information becomes available. Implementation of this strategy is subject to
30 appropriations, priorities and budgetary constraints of the participating jurisdictions and
31 organizations.

32 Success in the recovery of this species depends on the commitment and cooperation of
33 many different constituencies that will be involved in implementing the advice set out in
34 this strategy.

35 **Responsible jurisdictions**

36 Ministry of the Environment, Conservation and Parks
37 Environment and Climate Change Canada – Canadian Wildlife Service, Ontario
38 Parks Canada Agency
39

40 **Executive summary**

41 The *Endangered Species Act, 2007* (ESA) requires the Minister of the Environment,
42 Conservation and Parks to ensure recovery strategies are prepared for all species listed
43 as endangered or threatened on the Species at Risk in Ontario (SARO) List. Under the
44 ESA, a recovery strategy may incorporate all or part of an existing plan that relates to
45 the species.

46 The Eastern Whip-poor-will (*Antrostomus vociferus*) is listed as threatened on the
47 SARO List. The species is listed as threatened under the federal *Species at Risk Act*
48 (SARA). Environment and Climate Change Canada prepared the Recovery Strategy for
49 the Eastern Whip-poor-will (*Antrostomus vociferus*) in Canada in 2018 to meet its
50 requirements under the SARA. This recovery strategy is hereby adopted under the
51 ESA. With the additions indicated below, the enclosed strategy meets all the content
52 requirements outlined in the ESA.

53 Additions include new information on potential Eastern Whip-poor-will occurrences, and
54 the threat posed to this species by declines in aerial insect populations and loss of
55 habitat in migratory stopover areas.

56 The Critical Habitat section of the federal recovery strategy provides an identification of
57 critical habitat (as defined under the SARA). Identification of critical habitat is not a
58 component of a recovery strategy prepared under the ESA. However, it is
59 recommended that the approach used to identify critical habitat in the federal recovery
60 strategy, along with data available at the Ministry of Natural Resources and Forestry's
61 Natural Heritage Information Centre and any new scientific information concerning
62 Eastern Whip-poor-will and the areas it occupies, be considered when developing a
63 habitat regulation under the ESA.

64

65 **Table of contents**

66 Recommended citation..... i
67 Declarationii
68 Responsible jurisdictions.....ii
69 Executive summary.....iii
70 1.0 Adoption of federal recovery strategy 1
71 1.1 Species assessment and classification 1
72 1.2 Distribution, abundance and population trends 1
73 1.3 Threats to survival and recovery 1
74 1.4 Area for consideration in developing a habitat regulation 2
75 Glossary 3
76 List of abbreviations 3
77 References..... 5
78 Appendix 1. Recovery strategy for the Eastern Whip-poor-will (*Antrostomus vociferus*)
79 in Canada 6
80

81 **1.0 Adoption of federal recovery strategy**

82 The *Endangered Species Act, 2007* (ESA) requires the Minister of the Environment,
83 Conservation and Parks to ensure recovery strategies are prepared for all species listed
84 as endangered or threatened on the Species at Risk in Ontario (SARO) List. Under the
85 ESA, a recovery strategy may incorporate all or part of an existing plan that relates to
86 the species.

87 The Eastern Whip-poor-will (*Antrostomus vociferus*) is listed as threatened on the
88 SARO List. The species is listed as threatened under the federal *Species at Risk Act*
89 (SARA). Environment and Climate Change Canada prepared the Recovery Strategy for
90 the Eastern Whip-poor-will (*Antrostomus vociferus*) in Canada in 2018 to meet its
91 requirements under the SARA. This recovery strategy is hereby adopted under the
92 ESA. With the additions indicated below, the enclosed strategy meets all the content
93 requirements outlined in the ESA.

94 **1.1 Species assessment and classification**

95 The following list is a summary of assessment and classification information for the
96 Eastern Whip-poor-will (*Antrostomus vociferus*). Note: The glossary provides definitions
97 for the abbreviations and technical terms in this document.

- 98 • SARO List Classification: Threatened
- 99 • SARO List History: Threatened (2009)
- 100 • COSEWIC Assessment History: Threatened (2009)
- 101 • SARA Schedule 1: Threatened
- 102 • Conservation Status Rankings: G-rank: G5; N-rank: N4B, N3M; S-rank: S4B

103 **1.2 Distribution, abundance and population trends**

104 The federal recovery strategy summarizes Atlas of the Breeding Birds of Ontario data
105 (2001-2005) and North American Breeding Bird Survey Ontario data (2002-2012) to
106 characterize Eastern Whip-poor-will population trends in the province (Environment and
107 Climate Change Canada 2018). From 2013 to 2018, the Ministry of Natural Resources
108 and Forestry's (MNR) Natural Heritage Information Centre received close to 4000
109 observation reports of Eastern Whip-poor-will, a proportion of which are expected to
110 result in the identification of new verified occurrences (MNR 2018).

111 **1.3 Threats to survival and recovery**

112 Environment and Climate Change Canada (ECCC) (2018) identifies the reduced
113 availability of insect prey as the principal threat to Eastern Whip-poor-will. Analysis

114 conducted at multiple spatial scales suggests that food availability is an important
115 predictor of Eastern Whip-poor-will abundance in Ontario (English et al. 2017a), such
116 that the species' presence was significantly related to moth abundance at a regional
117 scale, and at a local scale its abundance was found to be significantly related to beetle
118 abundance.

119 ECCC (2018) identifies deforestation on the wintering grounds of Eastern Whip-poor-will
120 as the second greatest threat to the species. English et al. (2017b) also found evidence
121 that migratory stopover areas on the north coast of the Gulf of Mexico are important for
122 successful fall migration, where loss of forest cover may also be a concern.

123 **1.4 Area for consideration in developing a habitat regulation**

124 Under the ESA, a recovery strategy must include a recommendation to the Minister of
125 the Environment, Conservation and Parks on the area that should be considered in
126 developing a habitat regulation. A habitat regulation is a legal instrument that prescribes
127 an area that will be protected as the habitat of the species. The recommendation
128 provided below will be one of many sources considered by the Minister, including
129 information that may become newly available following completion of the recovery
130 strategy, when developing the habitat regulation for this species.

131 The Critical Habitat section of the federal recovery strategy provides an identification of
132 critical habitat (as defined under the SARA). Identification of critical habitat is not a
133 component of a recovery strategy prepared under the ESA. However, it is
134 recommended that the approach used to identify critical habitat in the federal recovery
135 strategy, along with observation data available at MNRF's Natural Heritage Information
136 Centre and any new scientific information on Eastern Whip-poor-will and the areas it
137 occupies, be considered when developing a habitat regulation under the ESA.

138 **Glossary**

139 Committee on the Status of Endangered Wildlife in Canada (COSEWIC): The
140 committee established under section 14 of the Species at Risk Act that is
141 responsible for assessing and classifying species at risk in Canada.

142 Committee on the Status of Species at Risk in Ontario (COSSARO): The committee
143 established under section 3 of the *Endangered Species Act, 2007* that is
144 responsible for assessing and classifying species at risk in Ontario.

145 Conservation status rank: A rank assigned to a species or ecological community that
146 primarily conveys the degree of rarity of the species or community at the global
147 (G), national (N) or subnational (S) level. These ranks, termed G-rank, N-rank
148 and S-rank, are not legal designations. Ranks are determined by NatureServe
149 and, in the case of Ontario's S-rank, by Ontario's Natural Heritage Information
150 Centre. The conservation status of a species or ecosystem is designated by a
151 number from 1 to 5, preceded by the letter G, N or S reflecting the appropriate
152 geographic scale of the assessment. The numbers mean the following:

- 153 1 = critically imperilled
- 154 2 = imperilled
- 155 3 = vulnerable
- 156 4 = apparently secure
- 157 5 = secure
- 158 NR = not yet ranked

159 *Endangered Species Act, 2007* (ESA): The provincial legislation that provides protection
160 to species at risk in Ontario.

161 *Species at Risk Act* (SARA): The federal legislation that provides protection to species
162 at risk in Canada. This Act establishes Schedule 1 as the legal list of wildlife
163 species at risk. Schedules 2 and 3 contain lists of species that at the time the Act
164 came into force needed to be reassessed. After species on Schedule 2 and 3 are
165 reassessed and found to be at risk, they undergo the SARA listing process to be
166 included in Schedule 1.

167 Species at Risk in Ontario (SARO) List: The regulation made under section 7 of the
168 *Endangered Species Act, 2007* that provides the official status classification of
169 species at risk in Ontario. This list was first published in 2004 as a policy and
170 became a regulation in 2008.

171 **List of abbreviations**

- 172 COSEWIC: Committee on the Status of Endangered Wildlife in Canada
- 173 COSSARO: Committee on the Status of Species at Risk in Ontario
- 174 ECCC: Environment and Climate Change Canada

175 ESA: Ontario's *Endangered Species Act, 2007*
176 ISBN: International Standard Book Number
177 MNRF: Ministry of Natural Resources and Forestry
178 SARA: Canada's *Species at Risk Act*
179 SARO List: Species at Risk in Ontario List
180

181

182 **References**

183 English, P.A., J.J. Nocera, B.A. Pond and D.J. Green. 2017a. Habitat and food supply
184 across multiple spatial scales influence the distribution and abundance of a
185 nocturnal aerial insectivore. *Landscape Ecology* 32:343-359.

186 English, P.A., A.M. Mills, M.D. Cadman, A.E. Heagy, G.J. Rand, D.J. Green and
187 J.J. Nocera. 2017b. Tracking the migration of a nocturnal aerial insectivore in the
188 Americas. *BMC Zoology* 2(5):1-11.

189 Environment and Climate Change Canada (ECCC). 2018. Recovery Strategy for the
190 Eastern Whip-poor-will (*Antrostomus vociferus*) in Canada. Species at Risk Act
191 Recovery Strategy Series. Environment and Climate Change Canada, Ottawa.

192 Ministry of Natural Resources and Forestry (MNRF). 2018. Provincially Tracked Species
193 Observations for Eastern Whip-poor-will (*Antrostomus vociferous*). Natural
194 Heritage Information Centre, Ministry of Natural Resources and Forestry,
195 Peterborough, Ontario. Accessed 29 November 2018.

196

197

198

**Appendix 1. Recovery strategy for the
Eastern Whip-poor-will (*Antrostomus vociferus*)
in Canada**

Recovery Strategy for the Eastern Whip-poor-will (*Antrostomus vociferus*) in Canada

Eastern Whip-poor-will



2018



Government
of Canada

Gouvernement
du Canada

Canada

Recommended citation:

Environment and Climate Change Canada. 2018. Recovery Strategy for the Eastern Whip-poor-will (*Antrostomus vociferus*) in Canada. *Species at Risk Act Recovery Strategy Series*. Environment and Climate Change Canada, Ottawa. vi + 107 pp.

For copies of the recovery strategy, or for additional information on species at risk, including the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) Status Reports, residence descriptions, action plans, and other related recovery documents, please visit the [Species at Risk \(SAR\) Public Registry](#)¹.

Cover illustration: © H. Loney Dickson

Également disponible sous le titre
« Programme de rétablissement de l'Engoulevent bois-pourri (*Antrostomus vociferus*)
au Canada »

© Her Majesty the Queen in Right of Canada, represented by the Minister of Environment and Climate Change, 2018. All rights reserved.

ISBN 978-0-660-28364-7

Catalogue no. En3-4/300-2018E-PDF

Content (excluding the illustrations) may be used without permission, with appropriate credit to the source.

¹ <http://sararegistry.gc.ca/default.asp?lang=En&n=24F7211B-1>

Preface

The federal, provincial, and territorial government signatories under the [Accord for the Protection of Species at Risk \(1996\)](#)² agreed to establish complementary legislation and programs that provide for effective protection of species at risk throughout Canada. Under the *Species at Risk Act* (S.C. 2002, c.29) (SARA), the federal competent ministers are responsible for the preparation of recovery strategies for listed Extirpated, Endangered, and Threatened species and are required to report on progress within five years after the publication of the final document on the SAR Public Registry.

The Minister of Environment and Climate Change and Minister responsible for the Parks Canada agency is the competent minister under SARA for the Eastern Whip-poor-will and has prepared this recovery strategy, as per section 37 of SARA. To the extent possible, it has been prepared in cooperation with the Provinces of Saskatchewan, Manitoba, Ontario (Ministry of Natural Resources and Forestry), Quebec (Ministère des Forêts, de la Faune et des Parcs), New Brunswick and Nova Scotia as per section 39(1) of SARA.

Success in the recovery of this species depends on the commitment and cooperation of many different constituencies that will be involved in implementing the directions set out in this strategy and will not be achieved by Environment and Climate Change Canada and the Parks Canada Agency, or any other jurisdiction alone. All Canadians are invited to join in supporting and implementing this strategy for the benefit of the Eastern Whip-poor-will and the Canadian society as a whole.

This recovery strategy will be followed by one or more action plans that will provide information on recovery measures to be taken by Environment and Climate Change Canada, the Parks Canada Agency, and other jurisdictions and/or organizations involved in the conservation of the species. Implementation of this strategy is subject to appropriations, priorities, and budgetary constraints of the participating jurisdictions and organizations.

The recovery strategy sets the strategic direction to arrest or reverse the decline of the species, including identification of critical habitat to the extent possible. It provides all Canadians with information to help take action on species conservation. When critical habitat is identified, either in a recovery strategy or an action plan, SARA requires that critical habitat then be protected.

In the case of critical habitat identified for terrestrial species including migratory birds SARA requires that critical habitat identified in a federally protected area³ be described

² <http://registrelep-sararegistry.gc.ca/default.asp?lang=en&n=6B319869-1#2>

³ These federally protected areas are: a national park of Canada named and described in Schedule 1 to the *Canada National Parks Act*, The Rouge National Park established by the *Rouge National Urban Park Act*, a marine protected area under the *Oceans Act*, a migratory bird sanctuary under the *Migratory Birds Convention Act, 1994* or a national wildlife area under the *Canada Wildlife Act* see ss. 58(2) of SARA.

in the *Canada Gazette* within 90 days after the recovery strategy or action plan that identified the critical habitat is included in the public registry. A prohibition against destruction of critical habitat under ss. 58(1) will apply 90 days after the description of the critical habitat is published in the *Canada Gazette*.

For critical habitat located on other federal lands, the competent minister must either make a statement on existing legal protection or make an order so that the prohibition against destruction of critical habitat applies.

If the critical habitat for a migratory bird is not within a federal protected area and is not on federal land, within the exclusive economic zone or on the continental shelf of Canada, the prohibition against destruction can only apply to those portions of the critical habitat that are habitat to which the *Migratory Birds Convention Act, 1994* applies as per SARA ss. 58(5.1) and ss. 58(5.2).

For any part of critical habitat located on non-federal lands, if the competent minister forms the opinion that any portion of critical habitat is not protected by provisions in or measures under SARA or other Acts of Parliament, or the laws of the province or territory, SARA requires that the Minister recommend that the Governor in Council make an order to prohibit destruction of critical habitat. The discretion to protect critical habitat on non-federal lands that is not otherwise protected rests with the Governor in Council.

Acknowledgments

This recovery strategy was prepared by Vincent Carignan (Environment and Climate Change Canada, Canadian Wildlife Service [ECCC-CWS]– Quebec region) and Mireille Poulin⁴ based on an initial draft by Kimberley Hair and Madison Wikston (ECCC-CWS- National Capital region) and in collaboration with:

ECCC-CWS - Ontario region : Ken Tuininga, Mike Cadman, Megan Eplett, Krista Holmes, Marie-Claude Archambault, Lyle Friesen, Jeff Robinson, Russ Weeber, Rich Russell, Kari Van Allen, Madeline Austen, Elizabeth Rezek and Lesley Dunn.

ECCC-CWS - Atlantic region: Julie McKnight, Peter Thomas, Krista Baker, David Andrews, Andrew Boyne and Becky Whittham.

ECCC-CWS - National Capital region: Manon Dubé and Adam Smith.

ECCC-CWS - Quebec region: Bruno Drolet, Gilles Falardeau, François Shaffer and Geneviève Langlois.

ECCC-CWS - Prairie & Northern region: Mark Bidwell, Steven Van Wilgenburg and John Conkin.

ECCC-CWS- Pacific & Yukon region: Pam Sinclair.

ECCC - Science & Technology: Junior Tremblay (Quebec region) et Scott Wilson (National Capital region).

Department of National Defence (DND): Robert Werbiski (Montreal/Saint-Jean garrison)

Parks Canada Agency (PCA): Joanne Tuckwell, Leah de Forest, Diane Amirault-Langlais and Eric Tremblay.

Ontario Ministry of Natural Resources and Forestry (OMNRF): Chris Risley, Aileen Rapson, Amanda Fracz and Jay Fitzsimmons (Species at Risk Branch); Erica Barkley and Jennifer Hoare (Ontario Parks); Joe Nocera (Wildlife Research and Monitoring Section); Southern Region: Jim Saunders, Brian Naylor (Forests Branch), Todd Norris (Peterborough District), Eric Cobb, Nikki Boucher (Northeast Region).

Additional thanks go to: Maureen Toner (New Brunswick Department of Natural Resources), Audrey Heagy (Bird Studies Canada – Ontario); Garry Gregory (Prince Edward Island Department of Communities, Land, and Environment) and Mark Elderkin (Nova Scotia Department of Natural Resources); Narayan Dhital (Forest Service, Saskatchewan); all other parties including Aboriginal organizations, landowners, citizens, and stakeholders who provided comments on the present document as well as Regroupement QuébecOiseaux, ECCC-CWS and Bird Studies Canada for providing the data from the Breeding Bird Atlases in Canada, and the thousands of volunteers that collected data for those projects.

⁴ Formerly with ECCC-CWS – Quebec region

Executive Summary

The Eastern Whip-poor-will (*Antrostomus vociferous*) is a nocturnal insectivorous bird that breeds in sparse forests or at the edge of forests adjacent to open habitats required for foraging. The species was designated as Threatened by the Committee for the Status of Endangered Wildlife in Canada (COSEWIC) in 2009 and has been listed under the same status in Schedule 1 of the *Species at Risk Act* (SARA) since 2011.

An estimated 120 000 Eastern Whip-poor-will individuals (6% of the global population) are found in Canada, where they breed in southern Saskatchewan, Manitoba, Ontario, Quebec, New Brunswick, Nova Scotia and Prince Edward Island. National and provincial population trends indicate a decline between 2.77% and 5.53% per year over the 2002-2012 period.

Although our understanding of the causes of the decline of the Eastern Whip-poor-will may be limited, the main threats include reduced availability of insect prey, agricultural expansion and intensification (wintering and breeding grounds), urban expansion, as well as energy development and mineral extraction. Other threats that contribute to a lesser extent are also presented.

There are unknowns regarding the feasibility of recovery of the Eastern Whip-poor-will. Nevertheless, in keeping with the precautionary principle, a recovery strategy has been prepared as per section 41(1) of SARA as would be done when recovery is determined to be feasible.

The population and distribution objectives for the Eastern Whip-poor-will in Canada are:

- In the short term: Slow the decline such that the population does not decrease by more than 10% (i.e. 12 000 individuals) over the 2018-2028 period, and maintain the area of occupancy at 3,000 km² or above;
- In the long term: Ensure a positive 10-year population trend starting in 2028, while favouring an increase in the area of occupancy, including the gradual recolonization of areas in the southern portion of the breeding distribution.

Broad strategies and approaches to achieve these objectives are presented in the Strategic Direction for Recovery section.

Critical habitat for the Eastern Whip-poor-will is partially identified in this recovery strategy. It corresponds to the areas of suitable nesting and/or foraging habitats within all 10 x 10 km standardized UTM squares with confirmed breeding or multiple occupancy since 2001. Overall, 198 critical habitat units are identified for the Eastern Whip-poor-will, including 32 in Manitoba, 110 in Ontario, 51 in Quebec, and five in the Maritimes (all in New Brunswick). A schedule of studies outlines the activities required to complete the identification of critical habitat.

One or more action plans will be posted on the Species at Risk Public Registry before the end of 2023.

Recovery Feasibility Summary

Based on the following four criteria that Environment and Climate Change Canada uses to establish recovery feasibility, there are unknowns regarding the feasibility of recovery of the Eastern Whip-poor-will. In keeping with the precautionary principle, this recovery strategy has been prepared as per section 41(1) of SARA, as would be done when recovery is determined to be technically and biologically feasible. This recovery strategy addresses the unknowns surrounding the feasibility of recovery.

1. Individuals of the wildlife species that are capable of reproduction are available now or in the foreseeable future to sustain the population or improve its abundance.

Yes. Breeding individuals are currently distributed in multiple areas of the Canadian breeding distribution as well as in the United States.

2. Sufficient suitable habitat is available to support the species or could be made available through habitat management or restoration.

Unknown. Sufficient suitable breeding habitat is available to support the species at its current level. Unoccupied and apparently suitable habitat appears to be available and additional habitats could become suitable after restoration efforts or through natural processes (e.g., succession, fires) or human activities (e.g., forest harvesting). The availability of sufficient amounts of migrating and wintering habitats is unknown.

3. The primary threats to the species or its habitat (including threats outside Canada) can be avoided or mitigated.

Unknown. Clarifying the impacts of some of the threats, namely the existence of possible thresholds in habitat loss as well as in the availability of insect prey populations would help establish more precise beneficial management practices. It is unknown whether threats such as reduced insect prey populations and threats to the wintering areas can be mitigated.

4. Recovery techniques exist to achieve the population and distribution objectives or can be expected to be developed within a reasonable timeframe.

Unknown. Habitat conservation and stewardship, along with habitat management techniques, could be effective for this species although specific management practices need to be developed and implemented. Mitigating threats, such as reduced insect prey populations and habitat availability on the wintering areas represent considerable challenges.

Table of Contents

Preface.....	i
Acknowledgments	iii
Executive Summary	iv
Recovery Feasibility Summary	v
1. COSEWIC Species Assessment Information.....	1
2. Species Status Information	1
3. Species Information	2
3.1 Species Description	2
3.2 Species Population and Distribution	3
3.3 Needs of the Eastern Whip-poor-will	5
4. Threats.....	8
4.1 Threat Assessment.....	8
4.2 Description of Threats.....	9
5. Population and Distribution Objectives	15
6. Broad Strategies and General Approaches to Meet Objectives	16
6.1 Actions Already Completed or Currently Underway	16
6.2 Strategic Direction for Recovery	18
6.3 Narrative to Support the Recovery Planning Table.....	21
7. Critical Habitat.....	22
7.1 Identification of the Species' Critical Habitat.....	22
7.2 Schedule of Studies to Identify Critical Habitat	26
7.3 Activities Likely to Result in the Destruction of Critical Habitat	27
8. Measuring Progress.....	29
9. Statement on Action Plans.....	30
10. References.....	30
Appendix A: Critical Habitat for the Eastern Whip-poor-will in Canada	42
Appendix B. Effects on the Environment and Other Species	107

1. COSEWIC* Species Assessment Information

Date of Assessment: April 2009

Common Name: Eastern Whip-poor-will

Scientific Name: *Antrostomus vociferus*

COSEWIC Status: Threatened

Reason for Designation: In Canada, this well-known, nocturnal bird has experienced both long-term and short-term population declines. Indices of abundance indicate that populations have been reduced by more than 30% over the last 10 years (i.e. 3 generations). Like other aerial foraging insectivores, habitat loss and degradation as well as changes to the insect prey base may have affected Canadian populations.

Canadian Occurrence: Saskatchewan, Manitoba, Ontario, Quebec, New Brunswick, Nova Scotia

COSEWIC Status History: Designated Threatened in April 2009.

* Committee on the Status of Endangered Wildlife in Canada. The information presented in this box, including the species' scientific name, reflects the state of knowledge at the time the species was evaluated.

2. Species Status Information

Approximately 6% of the global population and 20% of the breeding distribution of the Eastern Whip-poor-will (*Antrostomus vociferus*) is found in Canada (COSEWIC 2009; Partners in Flight Science Committee 2013). The species was listed as Threatened on Schedule 1 of the *Species at Risk Act* (SARA; S.C. 2002, c. 29) in April 2011. It is listed as Threatened under three provincial *Endangered Species Acts*: Ontario (S.O. 2007, c. 6), Nova Scotia (S.N.S. 1998, ch. 11), and Manitoba (C.C.C.S.M. c. E111, 1990), as well as under New Brunswick's *Species at Risk Act* (SNB 2012, c. 6). In Quebec, the species is listed as Likely to be designated as Threatened or vulnerable on the *List of Wildlife Species Likely to be Designated Threatened or Vulnerable*⁵ produced according to the *Act Respecting Threatened or Vulnerable Species* (CQLR, c. E-12.01). As of November 2014, the species was not listed in Saskatchewan nor Prince Edward Island.

NatureServe (2014) considers the global populations of the Eastern Whip-poor-will to be Secure (G5). The species' breeding populations are Apparently Secure (N4B) in Canada and Secure (N5B) in the United States. Table 1 provides further details on other conservation ranks in Canada.

⁵ With respect to the acquisition of knowledge, special attention is given to species on the *List of species likely to be designated threatened or vulnerable*.

Table 1. Conservation Status Ranks for the Breeding Populations of the Eastern Whip-poor-will in Canada (NatureServe 2014).

Region	Nature Serve ^a
Saskatchewan	Vulnerable (S3B)
Manitoba	Vulnerable (S3B)
Ontario	Apparently Secure (S4B)
Quebec	Vulnerable (S3)
New Brunswick	Imperiled (S2B)
Prince Edward Island	Non-breeder (SNA) ^b
Nova Scotia	Critically Imperiled (S1?B)

^a Conservation status ranks: NA: Not Applicable; ?: Inexact numerical rank; B: Breeding. For subnational ranks (by state) in the United States, refer to NatureServe's website <http://explorer.natureserve.org/>.

^b Note that some of these ranks have not been reviewed recently. For instance, both editions of the Breeding Bird Atlas in the Maritimes indicate that the species is present in at least one atlas square during the breeding season.

3. Species Information

The scientific name of the Eastern Whip-poor-will has been modified twice following the publication of the COSEWIC status report (2009). Initially, the Eastern Whip-poor-will (*Caprimulgus vociferus vociferus*) was considered a subspecies of the Whip-poor-will and encompassed all the eastern populations of the species' breeding distribution. In 2010, it was separated from its close relative, the Mexican Whip-poor-will (*Caprimulgus arizonae*), to form a new species by itself, the Eastern Whip-poor-will (*Caprimulgus vociferus*) (Chesser et al. 2010). In 2012, the *Antrorstomus* genus name, that was used for this species until 1931, was restored for the Eastern Whip-poor-will (refer to Chesser et al. 2012 for further details).

3.1 Species Description

Cink (2002) describes the Eastern Whip-poor-will as a nocturnal aerial insectivorous bird of the *Caprimulgidae* family (typical nightjars). It measures around 24 cm long and weighs 50 to 55 g. Its plumage is grey and brown, which serves to blend individuals with elements of the forest ground where they nest. Individuals have rounded wings and tail feathers and a large and flattened head with a small bill and big gape, bordered by long sensory bristles. Males differ from females in that they have a white collar and two big white patches on their outer tail feathers (smaller and buff-colored in females). Eastern Whip-poor-wills bear a striking resemblance to Common Nighthawks but lack the white patches on the wings and vocalizations are very distinct. As nocturnal birds, individuals are more often heard than seen and their distinct three-noted song "WHIP-poor-WEEL" is at the origin of the species' name.

3.2 Species Population and Distribution

The global breeding distribution of the Eastern Whip-poor-will is approximately 2 772 000 km² (COSEWIC 2009). It extends from Saskatchewan to the Maritimes and south in the United States, west to east from Oklahoma to Georgia (Cink 2002; Figure 1). The Eastern Whip-poor-will breeds locally throughout its range and no northward range expansion has been noted (Cink 2002). The global population is estimated at 2 million individuals (Partners in Flight Science Committee 2013). The Eastern Whip-poor-will's wintering range stretches from coastal South Carolina through Florida and along the Gulf Coast of the United States into Mexico and Central America as far south as Honduras and Panama (Cink 2002; Figure 1).

The Canadian breeding distribution occupies around 535 000 km² (20% of the global breeding distribution) (COSEWIC 2009) from east-central Saskatchewan, southern Manitoba, southern Ontario, southern Quebec, and sparse locations in New Brunswick, Nova Scotia and Prince Edward Island (Godfrey 1986; Cink 2002; Maritimes Breeding Bird Atlas 2013). The Canadian population is estimated at 120 000 individuals, with the highest concentrations found in eastern Ontario (Partners in Flight Science Committee 2013). Although this estimate incorporates atlas data, it is mainly based on data from the North American Breeding Bird Survey (BBS), and is considered to have a low reliability for the Eastern Whip-poor-will as this survey program is not designed to detect nocturnal birds (Partners in Flight Science Committee 2013; Environment Canada 2014a). Also, the Breeding Bird Survey data do not sample the species' entire range at random, having lower coverage in more remote areas. Nevertheless, Environment Canada's (2014a) analyses of the data indicate a high probability (0.86) that the population trend is declining.



Figure 1: Global range of the Eastern Whip-poor-will, which includes the breeding, wintering and migration distributions (Cornell Lab of Ornithology, 2007)

The Eastern Whip-poor-will's range within Canada has been retracting since the mid-1960s, most notably at its southern limit (Smith 1996; COSEWIC 2009). A comparison of the data from the first and second breeding bird atlases in Ontario and the Maritimes show a similar trend (Table 2). Although the atlas records have yet to be compiled for the last survey year of the second breeding bird atlas in Quebec, results indicate a slight increase in occupancy (Table 2). Differences in atlas square occupancy between successive atlases should be considered with caution as they can be the result of different levels of effort to detect the species.

Using BBS data, a Canada-wide population decline of 3.19% per year has been noted over the 1970-2012 period, or an approximate loss of 75% of the population (Environment Canada 2014a). Over the 2002-2012 period, this national trend has slightly slowed to a decline of 2.77% per year. Regional BBS trends for Ontario and Quebec, the core of the species' distribution in Canada, are consistent with the national trend (Table 2). Data from the Quebec check-list program (Étude des populations d'oiseaux du Québec - EPOQ) also corroborate these trends with a 2002-2011 population estimated to be about half that observed in the 1970's (Larivée 2013). It should be noted that declines are observed not only at the species level but also among

the guild of aerial insectivores and long-distance migrants to which the Eastern Whip-poor-will belongs (Blancher et al. 2009).

Table 2. Breeding Bird Atlases Data and Regional Breeding Bird Survey Trends.

Provinces	Atlas Periods	Number of Occupied Atlas Squares (trend 1 st vs 2 nd atlas)	BBS Annual Trends (1970-2012 / 2002-2012)	Atlas References
Saskatchewan	1966-2015 ^c	~36	Not available	Smith 1996; SBA
Manitoba	2010-2015	190	Not available	www.birdatlas.mb.ca
Ontario	1981-1985	885	-3.45% / -3.36%	Cadman et al. 1987
	2001-2005	554 (-37%)		Cadman et al. 2007
Quebec	1984-1989	168	-5.26% / -5.53%	Gauthier and Aubry 1996
	2010-2015 ^d	179 (+7%)		Atlas des oiseaux nicheurs du Québec 2014
Maritimes	1986-1990	62 (49 NB, 12 NS, 1 PE)	+0.53% / -0.09% (NB) ^e	Erskine 1992
	2006-2010	38 (-39%) (29 NB, 8 NS, 1 PE)		Maritimes Breeding Bird Atlas 2013

^c The Saskatchewan Bird Atlas (SBA) is not based on standard atlas methodology. Data are reported continually in a web-based application (gjsweb1.serm.gov.sk.ca/imf/imf.jsp?site=birds). Atlas squares correspond to the National Topographic System 1: 250 000 grids (nrcan.gc.ca/earth-sciences/geography/topographic-information/maps/9765) rather than the standard 10 x 10 km used in atlases.

^d Data reported is from the 2010-2013 period. Atlassing of the southern portion of the province was completed in 2015. Atlassing of the northern portion is still underway.

^e Trends are not available for the other provinces in the Maritimes due to the limited number of routes on which the species was detected.

3.3 Needs of the Eastern Whip-poor-will

Eastern Whip-poor-wills, like most forest birds, appear to rely on multi-scale considerations for habitat selection. Knowledge of the species' requirements at the landscape scale as well as during migrations and on the wintering grounds are currently limited.

Regional Context

The amount of forest cover, by providing more areas suitable for breeding, as well as the spatial configuration of forest habitats next to more open habitats are often reported as central to the species' presence (Roy and Bombardier 1996; Palmer-Ball 1996; Cink 2002, Wilson 2003; Garlapow 2007). Distance to larger forest tracts may also be important (Cink 2002), namely in more agricultural settings where the amount of nesting habitat is more limited.

Nesting and Foraging Habitats

The nesting and foraging habitats of the Eastern Whip-poor-will are usually defined by structural rather than compositional characteristics (Robbins 1994; Wilson and Watts 2008). Although the species is not considered particularly territorial, territory size ranges from 3 to 30 ha (Wilson 2003; Garlapow 2007; Hunt 2010) with a maximum value of 132.4 ha observed in central Ontario (mean 31 ha; Rand 2014). Home range size, the area within which birds are expected to be found most of the time as they conduct breeding activities, forage and move between these zones, can vary from 20 to 500 ha (mean 136 ha; Rand 2014). This is in line with Sandilands' (2010) estimation that 500 to 1000 ha may be necessary to support "more than a few pairs".

Nesting habitats

Nesting habitats include most types of forest at early stages of succession (or edges of forests with a dense tree cover but showing a similar structure at the ground level), rock or sand barrens with scattered trees, savannahs, old burns, as well as sparse conifer plantations (Wilson 1985; Bushman and Therres 1988; Cink, 2002; Mills 2007; Wilson and Watts 2008; Tozer et al. 2014). All these habitats exhibit characteristics such as well-drained soils, moderate tree cover (Godfrey 1986; Roy and Bombardier 1996; 26 to 83 % in Garlapow 2007) and moderate to sparse shrub and herbaceous cover (Eastman 1991; Garlapow 2007).

When woodlots are used for nesting (e.g., in agricultural landscapes), smaller isolated woodlots are not occupied by the species (Reese 1996), suggesting that there may be a threshold in forest patch size. However, Turpak et al. (2009) state that no threshold for size has been identified for the moment. Elevation also seems to play a role with the species usually being found at altitudes lower than 350 to 430 m (Cooper 1981; Robbins 1994; Roy and Bombardier 1996).

Although nests and eggs are very well camouflaged and rarely found, limited accounts show that eggs are laid directly on a thick bed of dead leaves (or bare ground), shaded by a short herbaceous plant, shrub or sapling (Roy and Bombardier 1996; Cink 2002), often near fallen tree limbs or rocks which can be used as perches to roost during the day (Cink 2002). Individuals appear to show a high degree of fidelity to nest sites (Cink 2002).

Foraging habitats

Foraging habitats include prairies, wetlands with shrubs, regenerating clearcuts as well as agricultural fields and other habitats with low tree cover and availability of foraging perches as these conditions favor the localisation of prey (lunar light) as well as foraging efficiency (Mills 1986; Garlapow 2007).

Foraging activities usually take place within 500 m of the nest, often near forest edges (Cink 2002; Garlapow 2007). However, at the northern edge of the species' distribution,

Rand (2014) showed significantly greater foraging distances that may result from reduced habitat quality and reduced flying insect abundance in regions where low temperatures hinder foraging activities.

Foraging takes place at dawn (up to 40 minutes after sunrise) or dusk (from 30 minutes after sunset until sufficient light remains available) (Mills 1986). Prey items consist mainly of large moths, scarab beetles and other nocturnal flying insects captured by sallying⁶ short distances from perches or picked directly on the ground (Tyler 1940; Cink 2002; Garlapow 2007).

Limiting Factors

Limiting factors influence a species' survival and reproduction, and play a major role in the capacity to attain certain population levels (recover following a decline). For the Eastern Whip-poor-will, these include:

- Low annual productivity: although the species may double-brood in some areas in southern Canada, it generally has a single small clutch consisting of two eggs (Peck and James 1983; Sandilands 2010).
- High predation: as a ground nester, the species may be particularly vulnerable to nest predation (Cink 2002; e.g. cats, coyotes, raccoons).

⁶ Sallying: To rush out or leap forth suddenly; a bird that sallies out from a branch to catch flying insects.

4. Threats

4.1 Threat Assessment

Table 3. Threat assessment for the Eastern Whip-poor-will.

Threat	Level of Concern ^f	Extent	Occurrence	Frequency	Severity ^g	Causal Certainty ^h
Habitat Loss or Degradation						
Agricultural expansion and intensification – wintering grounds	High	Widespread	Current	Continuous	Moderate	High
Agricultural expansion and intensification – breeding grounds	Medium	Localized	Current	Continuous	Moderate	High
Urban expansion	Medium	Localized	Current	Continuous	Moderate	Medium
Energy development and mineral extraction	Medium	Localized	Current	Continuous	Moderate	Medium
Overgrazing of forest understory	Low	Localized	Current	Continuous	Moderate	Medium
Forest management	Low	Widespread	Current	Continuous	Low	Low
Changes in Ecological Dynamics or Natural Processes						
Reduced availability of insect prey	High	Widespread	Current	Continuous	Unknown	Medium
Fire suppression	Low	Widespread	Current	Recurrent	Low	Low
Natural Processes or Activities						
Habitat succession	Medium	Localized	Current	Continuous	Moderate	Medium
Climate and Natural Disasters						
Climate change	Medium	Widespread	Current	Continuous	Unknown	Medium

^f *Level of Concern: signifies that managing the threat is of (high, medium or low) concern for the recovery of the species, consistent with the population and distribution objectives. This criterion considers the assessment of all the information in the table.*

^g *Severity: reflects the population-level effect (high: very large population-level effect, moderate, low, unknown).*

^h *Causal certainty: reflects the degree of evidence that is known for the threat (high: available evidence strongly links the threat to stresses on population viability; medium: there is a correlation between the threat and population viability, e.g., expert opinion; low: the threat is assumed or plausible).*

4.2 Description of Threats

Threats are listed in decreasing order of level of concern. It should be noted that there are knowledge gaps relating to the quantitative impacts of some of the threats, particularly during migrations and on the wintering grounds. There are also regional differences in the activities that are driving the declines of the species and its habitats (e.g., agricultural versus forestry landscapes). Secondary threats such as collisions (e.g., with vehicles along roads), predation, interspecific competition (Chuck-will's willow [*Antrostomus carolinensis*]) and invasive species (exotic earthworms) are not addressed directly in the present section but are considered in the Recovery planning table (section 6).

4.2.1. Reduced Availability of Insect Prey

The decline of aerial insectivores has been more obvious since the 1980s (Goldstein et al. 1999; Boettner et al. 2000; Wickramasinghe et al. 2004; Sauer et al. 1996; Blancher et al. 2007) and strongly suggests a cause related to changes in aerial insect populations on the breeding grounds, migration routes or wintering grounds (Nebel et al. 2010; Paquette et al. 2014). Indeed, insect populations are exhibiting significant declines worldwide, including in the United States (Laughlin and Kibbe 1985; Peterson and Meservey 2003). A recent review of global faunal population trends, noted that 33% of all insects with available IUCN-documented population trends were declining and many also exhibited range retractions (Dirzo et al. 2014). Declines are more severe in heavily disturbed regions, such as the tropics (Dirzo et al. 2014). In the northeastern United States, Wagner (2012) noted declines for many nocturnal moths, the preferred prey items for Eastern Whip-poor-will (Cink 2002).

The main possible causes for reduced availability of insect prey are identified and described below. Additional causes include changes in the predator communities, light proliferation, and acid precipitation (Graveland 1998; Benton et al. 2002; Wagner 2012; Goulson 2013). However, a direct causal effect between aerial insectivore abundance and insect populations has yet to be established, partly because long-term monitoring data on avian diets are limited (Nocera et al. 2012).

4.2.1.1. Loss of insect producing habitats

Many insects are limited to specific habitats for all or part of their life cycle and any activities that affect these habitats through habitat loss or intensification of uses (e.g., monocultures, pesticide and fertilizer-dependent crops; traditional versus conservation tillage; removal of hedgerows, wetland drainage, peat extraction, urban and coastal development; resource extraction; herbicides to control vegetation in utility rights-of-ways) would have impacts on their populations (U.S. Bureau of Land Management 1978, Foster 1991; Benton et al. 2002, Cunningham et al. 2004; Price et al. 2011, Brooks et al. 2012; Chiron et al. 2014).

Loss of insect producing habitats would affect Eastern Whip-poor-wills on the breeding grounds, along migration routes and on their wintering grounds.

4.2.1.2. Pesticides and other toxins

The use of pesticide in agriculture, forestry, and for mosquito control programs in urban areas has undoubtedly been a factor in the reduction of the abundance of flying insects throughout the Eastern Whip-poor-will's distribution.

Although most organochlorine pesticides (chemicals in the same family as dichlorodiphenyltrichloroethane – DDT) have been banned in North America for decades, there is indication that Neotropical migrant insectivores are still being exposed to their effects elsewhere in the Americas (Sager 1997, Klemens et al. 2000). These chemicals can have long-lasting effects on insect communities and thus the birds that rely on them. Dietary records of Chimney Swifts (*Chaetura pelagica*), an aerial insectivore, confirm a marked decrease in beetles (Coleoptera) and an increase in true bugs (Hemiptera) that was temporally correlated with a steep rise in DDT and its metabolites. Nocera et al. (2012) argued that DDT caused declines in Coleoptera and dramatic (possibly permanent) shifts in the insect communities, resulting in a nutrient-poor diet and ultimately a declining Chimney Swift population.

Non-selective pesticides, such as those used to control Gypsy moths or the more widely used neonicotinoids, have also been shown to affect aerial insectivores by reducing their prey populations and impairing reproduction (Colburn et al. 1993, Wickramasinghe et al. 2004; Mineau and Palmer 2013; Hallmann et al. 2014). Neonicotinoids are generally used in agricultural habitats, but have been detected in wetlands (Main et al. 2014) and waterways in Canada (Environment Canada 2011a, Xing et al. 2013). In 2013, the European Food Safety Authority declared that they posed an ‘unacceptable’ risk to insects (Goulson 2014). Mineau and Palmer (2013) suggested that the effects of neonicotinoids on birds may not be limited to the farm scale, but likely expand to the watershed or regional scale.

Although not necessarily related to a decline in flying insect populations, the consumption of mercury-contaminated insects has been shown to decrease reproductive success, alter immune responsiveness, and cause behavioural and physiological effects in many insectivorous bird species (Scheuhammer et al. 2007, Hawley et al. 2009). Increased mercury levels may result from multiple causes (e.g., acid depositions, creation of reservoirs) and many terrestrial songbirds of northeastern North America that eat invertebrates have been found to biomagnify the substance to level that may be of conservation concern (Osborne et al. 2011; Keller et al. (2014). Although the Eastern Whip-poor-will was not part of these studies, the conclusions likely apply to the species.

4.2.1.3. Insect/breeding temporal mismatch

Birds often exhibit a strong synchronization between their reproductive timing (i.e. hatching) and peak food abundance, but climate change has caused the timing of peaks in some insects to occur sooner in the season (Both et al. 2009). Because warming may be less severe on the Eastern Whip-poor-will's wintering grounds than on their breeding grounds, they may experience migration cues at dates that are too late for them to arrive at breeding grounds at the optimal time (Jones and Cresswell 2010). As a result, climate change is creating a temporal mismatch between reproduction and maximal prey abundance (i.e. insects) for species that are not adapting to the changing climate at the same rate as their prey (Strode 2003; Both et al. 2006; Gornish and Tylianakis, 2013). This has been shown to affect the weight of Great Tit (*Parus major*) chicks and the number of chicks that fledge (Visser et al. 2006). An insect/breeding temporal mismatch has also been linked to the population declines of migrant birds across Europe (Møller et al. 2008, Saino et al. 2011).

Long-distance migratory birds that breed in habitats where the peaks of food abundance are shorter (such as forests) are more vulnerable to climate change because the temporal mismatch is more likely and more severe (Both et al. 2006, Both et al. 2009). Although no species-specific data are currently available, the Eastern Whip-poor-will is a long-distance migratory aerial insectivore that breeds / forages in habitats where peaks of food abundance are shorter (usually late Spring/early Summer), so a climate-induced mismatch between breeding and prey availability is probable.

4.2.2. Agricultural Expansion and Intensification – wintering grounds

On the wintering grounds in Mexico and Central America, since the early 1900s, forest conversion into pastures for cattle ranching through slash-and-burn agriculture has driven the majority of deforestation (Leonard 1987; Masek et al. 2011; Aide et al. 2013). In that region, the countries that experienced the most forest area loss between 2001 and 2010 are Guatemala and Nicaragua, representing 3 019 km² and 7 961 km² respectively (Aide et al. 2013) but most countries exhibit substantial declines in forest cover (Hansen et al. 2013). Although Eastern Whip-poor-wills may benefit from habitat created by some types of conversion for cattle pastures, intensive conversion that eliminates corridors and leaves little to no forested habitat adjacent to open habitats used for foraging decreases habitat suitability.

The impact of this threat on the decline of the Eastern Whip-poor-will has not been quantified, but it is viewed as a concern because of the species' concentration in a limited amount of winter habitat.

4.2.3. Agricultural Expansion and Intensification – breeding grounds

In Canada, the increased demand for agricultural land was responsible for approximately 43% of the deforestation in 2010 (~20 000 ha), mainly in Alberta,

Saskatchewan, and Manitoba where the boreal forest borders the prairies⁷. In the eastern United States, rates of conversion of forest to agricultural land have slowed considerably in recent years (Ramankutty et al. 2010; Masek et al. 2011) and many areas are returning to a forested state.

Efforts to increase the productivity of agricultural lands have led to the loss of large natural habitats, loss of habitat diversity in agricultural landscapes (e.g. conversion to row crops, elimination of hedges and other natural features), and an increase in pollution from nutrients and pesticides used on crops, all of which affect biodiversity (Robinson and Sutherland 2002). Insectivorous bird abundance and richness are higher in agricultural landscapes that provide a heterogeneous array of habitats (Parish et al. 1995; Jones et al. 2005), which are also home to more insect prey (Lewis and Stephenson 1966; Lewis and Dibley 1970; Verboom and Spoelstra 1999). Furthermore, landscapes that are highly simplified (e.g., dominated by hard edges between intensive crops and adjacent forests) may not provide enough suitable habitat for the Eastern Whip-poor-will. In Quebec, it has been noticed that the species avoids large areas of intensive agriculture (Cyr and Larivée 1995; Bélanger et al. 1999). In southern Ontario, the conversion of open grasslands and thickets to intensive agriculture has also been identified as a threat to the species (Ontario Ministry of Natural Resources 2009).

4.2.4. Urban Expansion

Urban development is an increasing threat that leads to permanent habitat losses both in suburban and rural areas. It is considered the leading cause of deforestation in the United States, as well as a major contributing factor in Canada (17%)(Robinson et al. 2005; Radeloff et al. 2005; Sun et al. 2007; Masek et al. 2011; Jobin et al. 2014). In Pennsylvania (USA), urbanization, through the increase of predation and the decrease in availability of nesting and foraging habitat, is presumed responsible for the decline of the Eastern Whip-poor-will (Santner 1992).

Habitat loss and fragmentation by urbanization may also increase water, air, noise and light pollution. Ecological light pollution has been proven to disrupt predator-prey dynamics, competition dynamics, nest site choice, orientation and communication for a variety of animals (Longcore and Rich 2004). It could also have positive components (e.g., better foraging efficiency).

4.2.5. Energy Development and Mineral Extraction

Exploration to find new energy sources (e.g., oil, gas, coal and hydroelectricity) and minerals (including aggregates), exploitation of these sources (e.g., windfarms, mine residues, flooding of areas to create reservoirs) and their transportation (e.g., pipelines, transmission lines, roads) have generated habitat loss, degradation and fragmentation in some areas of the Eastern Whip-poor-will's distribution (Masek et al. 2011). Activities associated with these industries can also lead to the unintentional destruction of nest, eggs, nestling, and/or adults (Van Wilgenburg et al. 2013).

⁷ <http://www.nrcan.gc.ca/forests/inventory/13419>

Rand (2014) found that Eastern Whip-poor-wills did not show elevated stress levels (measured by corticosterone) from exposure to mining exploration activities. Otherwise, quantitative effects of the more active exploitation phase of mining on Eastern Whip-poor-will populations have not been measured, but are presumed to be important given that habitat alterations by these activities are often permanent.

The maintenance of some infrastructures used to transport energy can generate habitat that will eventually be suitable for foraging.

4.2.6. Overgrazing of forest understory

Forests overgrazed by cattle are usually avoided for nesting purposes by the Eastern Whip-poor-will (Tyler 1940; Godfrey 1986), although they may be used for foraging if suitable nesting habitat is nearby (Cink 2002). Deer grazing can also be problematic in some areas as their populations have been increasing for decades (Russell et al. 2001). In Canada, cattle overgrazing has been identified as a factor contributing to the disappearance of the species, particularly in south-eastern Saskatchewan's aspen forests (Smith 1996; Jorgenson and Foster 2007).

Whatever the source, overgrazing of forest understory decreases vegetation cover and modifies plant community composition and dynamics (Patric and Helvey 1986; Côté et al. 2004; Rooney 2009; Goetsch et al. 2011), which in turn negatively affects wildlife by altering nest cover and food sources (Gallizioli 1979; DeGraaf et al. 1991; Pollard and Cooke 1994; Gill 2000; Allombert et al. 2005). At present, it is unknown how this threat is affecting the Eastern Whip-poor-will's population, but it would be prominent in northeastern United States and south-eastern Canada where White-tailed Deer (*Odocoileus virginianus*) are particularly abundant (Russell et al. 2001).

4.2.7. Habitat Succession

In forested landscapes, the Eastern Whip-poor-will often takes advantage of the open areas created by low-intensity agriculture or forest management for foraging, while relying on adjacent forests for nesting (COSEWIC 2009). Agricultural land abandonment creates early- and mid-successional forests that can, at first, provide suitable habitat for the species, but succession eventually leads to older forest stages, which are not preferred habitats (Bushman and Therres 1988).

Although cropland areas have increased over the past 50 years in most of the Eastern Whip-poor-will's range in southern Canada, succession of abandoned farmland has been the trend in marginal areas (Cadman et al. 2007; Latendresse et al. 2008), such as eastern Ontario (Ontario Ministry of Natural Resources 2009) and the eastern United States where this succession has been implicated in the Eastern Whip-poor-wills' decline (Medler 2008). In those areas, forest succession may have caused some degree of nesting/foraging habitat loss for the Eastern Whip-poor-will (Mills 2007; Smith 1996).

4.2.8. Climate Change

The potential effects of climate change on the Eastern Whip-poor-will can be difficult to predict because different bird species respond differently to spatial and temporal variations in their environment (Taper et al. 1995). One effect, the increase in the number of severe weather events (cold snaps, hurricanes, wind storms), can have an impacts on the breeding grounds, along the migration routes and on the wintering grounds. Cold, wet weather during the breeding season is well known for affecting aerial insectivores (e.g., Brown and Brown 2000). Such fluctuations and weather extremes are expected to occur more frequently due to climate change (Huber and Gulledge 2011).

Fire activity is also strongly influenced by weather (Flannigan et al. 2009) and the extent, intensity, and frequency of forest fires are projected to increase due to warmer springs and summers and decreases in water availability in some areas (Flannigan et al. 2009, North American Bird Conservation Initiative US Committee 2010, de Groot et al. 2013, Girardin et al. 2013). This would have a positive effect on Eastern Whip-poor-will habitat availability.

During the Fall migration, tropical storms and hurricanes can kill individuals in large numbers; a single hurricane (Hurricane Wilma in 2005) led to a decline of 50% in the population of the Chimney Swift in Québec (Dionne et al. 2008). On the tropical and subtropical wintering areas, Anodon et al. (2014) suggest that climate change will result in the expansion of savannahs at the expense of forests, which could eventually affect Eastern Whip-poor-wills.

4.2.9. Fire Suppression

Among multiple habitat types used by the Eastern Whip-poor-will, the species may be associated with those created by fires (Cink 2002). Fire in a natural system may provide a shifting mosaic of nesting and foraging habitat. Modern fire suppression may keep forest stands into a more mature stage, less suitable for the species. Over smaller expanses, fire suppression has been identified as one of the causes of the bird's decline in Ontario (Mills 2007; Tozer et al. 2014). The extent of this threat is not quantified for Eastern Whip-poor-will populations.

4.2.10. Forest Management

Between 2000 and 2012, approximately 11 million ha of forest were harvested throughout Canada (NFD 2014). Forest harvesting rates in most of Canada have been relatively stable throughout the past 20 years but are expected to decline in the coming years (Masek et al. 2011).

Forest harvesting can have short term negative effects on nesting birds by disrupting breeding activities (Hobson et al. 2013). The nests and/or eggs can be inadvertently harmed or disturbed as a result of clearing trees and other vegetation

(e.g. pre-commercial thinning) (Environment Canada 2014b). Nesting failure could also result from disruptive activities experienced by a nesting bird (Environment Canada 2014b). Hobson et al. (2013) estimated that between 616,000 and 2.09 million nests (of many species) are lost annually as a result of industrial forest harvesting.

However, forest management can also improve habitat through practices such as clearcut interspersion with mature forests (Wilson and Watts 2008, Tozer et al. 2014), variable density thinning, early thinning and other aspects of partial cutting (Bushman and Therres 1988).

Much of the northern distribution of the Eastern Whip-poor-will in Canada is under forest management. Forest harvesting is also a prevalent activity along the migratory routes and on the wintering grounds.

5. Population and Distribution Objectives

The population and distribution objectives for the Eastern Whip-poor-will in Canada are:

- In the short term: Slow the decline such that the population does not decrease by more than 10% (i.e. 12 000 individuals) over the 2018-2028 period, and maintain the area of occupancy at 3,000 km² or above.
- In the long term: Ensure a positive 10-year population trend starting in 2028, while favouring an increase in the area of occupancy, including the gradual recolonization of areas in the southern portion of the breeding range.

These objectives address the species' long-term decline, which was the reason for its designation as Threatened (COSEWIC 2009). The 10-year time frame for the short term objectives corresponds to the period between successive COSEWIC assessments of a species status and is considered reasonable given the challenge halting the Eastern Whip-poor-will's decline represents. The area of occupancy provided corresponds to the COSEWIC (2009) estimate (33 000 Eastern Whip-poor-will pairs occupying 1,650 km²) but with the updated Canadian population (120 000 individuals or 60 000 pairs) provided by the Partners in Flight Science Committee (2013), for a total of 3,000 km². As for the long term objectives, due to the substantial loss of suitable habitat in the southern portion of the species' range (agricultural and urban landscapes), it may be unrealistic to bring Eastern Whip-poor-will populations back to their historical levels. Nevertheless, ensuring that suitable habitat is available throughout the species' range in Canada, including through prevention of degradation or loss or through restoration of habitats in highly transformed landscapes, is necessary for the management and recovery of the Eastern Whip-poor-will.

These objectives may be reviewed during the development of the report required five years after this strategy is posted to assess the implementation of the strategy and the progress towards meeting its objectives (s. 46 SARA).

6. Broad Strategies and General Approaches to Meet Objectives

6.1 Actions Already Completed or Currently Underway

Numerous activities have been initiated since the latest COSEWIC assessment (2009). The following list is not exhaustive, but is meant to illustrate the main areas where work is already underway, to give context to the broad strategies outlined in section 6.2.

- Multiple projects targeting the Eastern Whip-poor-will on federal, provincial and private lands with funding from the Habitat Stewardship Program, the Interdepartmental Recovery Fund and the Aboriginal Fund for Species at Risk.

Manitoba

- Surveys on Department of National Defence establishments.

Saskatchewan

- Research underway on the broad-scale predictors of aerial insectivorous bird declines across North America (Dr. Christy Morrissey's Ecotoxicology Lab at the University of Saskatchewan <http://homepage.usask.ca/~cam202/page11.html>).

Ontario

- General Habitat Description completed under the Endangered Species Act, 2007 <http://www.ontario.ca/environment-and-energy/eastern-whip-poor-will>
- Roadside surveys conducted by Bird Studies Canada (2010-2013) and provincial partners such as Ministry of Natural Resources and Forestry in many districts and Ontario Parks (e.g., Algonquin Park). Local efforts are continuing.
- Geolocators deployed in four regions of Ontario (2011 and 2012) (Mike Cadman – EC-CWS Ontario; Philina English).
- Bird Studies Canada habitat use study underway (2013-2015) at site in southwestern Ontario.
- Study on habitat use, types of forest management and landscape characteristics is underway (John Vandebroek - OMNRF Fort Frances District; English and Conboy 2013).
- Surveys on Department of National Defence establishments.

Quebec

- Roadside surveys coordinated by Regroupement QuebecOiseaux on 25 permanent BBS-style routes in agricultural landscapes (2012-ongoing).
- Surveys along potential routes in the Outaouais region by Dendroica Environnement et Faune (2013 – ongoing).

- Project funded by the Interdepartmental Recovery Fund and the Department of National Defence to conduct surveys of the populations, describe the location of suitable habitat and propose potential mitigation measures for military activities at five Department of National Defence establishments (2012-ongoing).

Maritimes

- Habitat Stewardship Program projects on forest birds at risk to assist in the development of Maritimes-specific beneficial management practices conducted by Bird Studies Canada (ongoing).

6.2 Strategic Direction for Recovery

Table 4. Recovery Planning for the Eastern Whip-poor-will.

Threat or Limitation	Broad Strategy for Recovery	Priority ⁱ	General Description of Research and Management Approaches
All threats	Stewardship and management of the species and its suitable habitat	High	<ul style="list-style-type: none"> – Support habitat conservation using legal measures (where applicable), stewardship agreements and other regulatory tools on the breeding grounds, along migration routes and on the wintering grounds – Support Beneficial Management Practices (BMP) development and implementation at the local and landscape levels to increase population size as well as the areas of occupied habitat <ul style="list-style-type: none"> – BMPs should prioritize the importance of maintaining/creating insect producing habitats – Integrate BMPs for Eastern Whip-poor-will with BMPs for other wildlife within a heterogeneous and dynamic mosaic – Use management techniques over large land units and/or within an ecosystem approach – Restore habitats in some highly modified landscapes to promote the recolonisation of portions of the global distribution range

Threat or Limitation	Broad Strategy for Recovery	Priority ⁱ	General Description of Research and Management Approaches
Knowledge gaps	Monitoring and Research	High	<ul style="list-style-type: none"> - Develop and implement standardized research and monitoring protocols to gather information on the species' ecology, populations and trends, habitat characteristics and insect prey populations, including: <ul style="list-style-type: none"> - Determine habitat attributes (e.g., amount, spatial scale, composition and configuration), insect prey densities/composition and other variables that influence the selection and quality of nesting and foraging habitat - Clarify the drivers of spatial aggregation found in some areas and their contribution in population dynamics - Study links between insect prey populations (including trends), bird diets in habitats of various qualities and reasons for decline (including the effect of insect control programs) - Elucidate links between insect populations and migration phenology, especially in the context of climate change - Conduct electronic acoustic monitoring in less accessible parts of the range (e.g., southern portion of the Canadian Shield) - Establish dispersal distances and migratory paths - Determine if important migratory stopover sites exist for this species and proceed with their description (see Mehlman et al. 2005) - Clarify impacts of threats on the wintering areas and along migratory routes - Develop, validate or improve habitat models (e.g., Boreal Avian Modelling Initiative; Broeckaert 2012; Broeckaert and Bédard 2012) - Determine the effects of cats and other predators on this species
		Low	<ul style="list-style-type: none"> - Survey historical occurrences at regular intervals (e.g., 10 years)
All threats	Education and Partnerships	Medium	<ul style="list-style-type: none"> - Promote municipal, provincial, national and international cooperation and coordination of local and landscape-level planning (e.g. Bird Conservation Regions and Conservation Strategies^j) to mitigate threats on the breeding, migrating and wintering ranges - Examine the necessity for a working group for the species or for nocturnal aerial insectivores or species that require successional habitats - Establish partnerships with governmental departments and agencies, groups interested in the species, private owners and the general public for a local/regional/territorial implementation of the recovery strategy - Make all documentation relative to the species and its habitat readily available to the various stakeholders (e.g., websites, publications)

Threat or Limitation	Broad Strategy for Recovery	Priority ⁱ	General Description of Research and Management Approaches
All threats	Law and Policy	High	<ul style="list-style-type: none"> – Promote the compliance with existing environmental laws, regulations and policies, namely the <i>Migratory Birds Convention Act, 1994</i> and associated regulations to prevent disturbance to adults, nests and eggs for all types of activities and land tenure^k
		Medium	<ul style="list-style-type: none"> – Encourage the implementation of existing policies and reduction programs for pesticides, light pollution, greenhouse gasses and other pollutants; management of predator populations (where needed) and develop new policies and programs where gaps exist

ⁱ“Priority” reflects the degree to which the broad strategy contributes directly to the recovery of the species or is an essential precursor to an approach that contributes to the recovery of the species.

^j Environment and Climate Change Canada’s website on Bird Conservation Regions and Conservation Strategies
<https://www.ec.gc.ca/mbc-com/default.asp?lang=En&n=1D15657A-1>

^k Environment and Climate Change Canada’s website on the incidental take of migratory birds : <http://www.ec.gc.ca/paom-itmb/default.asp?lang=En&n=C51C415F-1>

6.3 Narrative to Support the Recovery Planning Table

Stewardship and Management of the Species and its Suitable Habitat

Preserving and enhancing Eastern Whip-poor-will nesting and foraging habitats will require promotion of approaches on a broad scale. These habitats must be managed and protected to ensure species survival, particularly in areas where large portions of habitat have been lost or degraded or are facing increased development pressures.

The key actions that can be promoted include beneficial management practices in forest management and agricultural activities as well as habitat restoration to increase habitat supply, where needed. Beneficial management practices for the Eastern Whip-poor-will must be integrated with those for other species to maintain heterogeneous landscapes that are a dynamic mosaic of habitat conditions which will benefit several species. Whenever possible, a multi-species or ecosystem approach to recovery should be considered. Beneficial management practices for governments, Aboriginal communities, industry, and even individuals can play an important role for the ongoing efforts to promote recovery of the Eastern Whip-poor-will. However, it must be kept in mind that such approaches will fail to recover the species unless migration and wintering habitats are also maintained and threats are addressed.

Monitoring and Research

A comprehensive approach to research and monitoring (that includes all stages of the annual life cycle and the entire range of occupancy) will be required to more completely understand the status of the species, as well as its threats and limiting factors on the breeding, migration and wintering grounds. Currently, adequate monitoring of the species is limited and concentrated close to urban centres. Targeted research and monitoring efforts are required throughout the Eastern Whip-poor-will's range in order to determine key demographic parameters (e.g., survival, dispersal, reproductive success in different habitat types) and identify migratory routes, stopover sites, and migratory connectivity that provide insight into the species needs at multiple scales. Trends in habitats and prey populations must be better understood (and monitored) to know whether maintaining, enhancing, and/or restoring insect producing habitats would significantly benefit Eastern Whip-poor-will populations.

Because the broad-scale monitoring of the Eastern Whip-poor-will is complicated by the large range of the species and its nocturnal habits, using alternative survey approaches (e.g., electronic acoustic monitoring) should be explored and citizen-based databases such as eBird should be considered. There are fewer monitoring programs established on the wintering grounds, but these are essential. They need to be developed and implemented to provide better information on habitat use, local habitat and population trends. Associated with these efforts, there is a need to build and validate corresponding habitat models at national and regional scales to better understand where on the landscape the species would be expected to breed, and assist with efforts to protect important and critical habitat.

Education and Partnerships

Collaboration with national and international jurisdictions (all levels of government), aboriginal peoples, industries, non-governmental organizations and landowners to preserve, restore, and enhance breeding, migration and wintering habitats is an important component of this recovery strategy. This could take the form of a working group such as those that have been put together for the Golden-winged Warbler (*Vermivora chrysoptera*) and the Canada Warbler (*Wilsonia canadensis*) or could target a larger group of species such as nocturnal aerial insectivores. Indeed, to be effective, research, conservation, and stewardship approaches must be applied throughout the species' range.

Promoting volunteer participation and collaborations in monitoring programs will also be a key element. Volunteers whose efforts should be promoted include Aboriginal communities (traditional ecological knowledge), local bird clubs who have knowledge of areas with high breeding densities and citizen scientists participating in bird atlases, breeding bird survey programs and citizen-based data collection (e.g., eBird).

Law and Policy

Throughout the species range, promoting the compliance with existing regulations and policies should be a priority. Currently, there are multiple legal means available to protect Eastern Whip-poor-wills and their habitats in Canada (e.g., Endangered Species Acts). It is important that these tools are fully realized and utilized for the protection of the species.

General prohibitions under the *Migratory Birds Convention Act, 1994* and its regulations also protect Eastern Whip-poor-will adults, young, nests and eggs anywhere they are found in Canada, regardless of land ownership. During the breeding period, potential destructive or disruptive activities should be avoided at locations where the species is likely to be encountered (Environment Canada 2014a).

7. Critical Habitat

7.1 Identification of the Species' Critical Habitat

SARA defines critical habitat as "...the habitat that is necessary for the survival or recovery of a listed wildlife species...". Section 41(1)(c) of SARA requires that recovery strategies include an identification of the species' critical habitat, to the extent possible, as well as examples of activities that are likely to result in its destruction. This recovery strategy partially identifies critical habitat, based on the best available information for the Eastern Whip-poor-will as of September 2014. The Schedule of Studies (section 7.2) outlines the activities required for completing the identification of the critical habitat necessary to meet the population and distribution objectives. As additional information becomes available, more precise boundaries may be mapped and further critical habitat may be identified.

The identification of critical habitat for the Eastern Whip-poor-will is based on two criteria: habitat occupancy and habitat suitability.

7.1.1 Habitat Occupancy

This criterion is intended to identify, with a reasonable degree of certainty, the areas of nesting and foraging habitats used by the species. In an analysis of a large number of studies of North American and European birds, Bock and Jones (2004) found that habitat occupancy was frequently an appropriate indicator of habitat suitability.

Habitat occupancy is based on documented nest locations, standardized survey data, as well as incidental observations. Confirmed breeding records^{8,9} constitute the highest indication of habitat occupancy. However, as the Eastern Whip-poor-will is a nocturnal species, breeding confirmation is rare. Accordingly, a more precautionary approach to establishing occupancy is warranted and considers combinations of records with lower nesting probabilities (probable or possible breeding birds).

Because Breeding Bird Atlases are the main sources of Eastern Whip-poor-will data across Canada, the earliest fieldwork year among the second editions of atlas projects (2001 in Ontario) was determined as the starting year to establish occupancy using all available data sources (species-specific surveys, BBS, Conservation Data Centres and check-lists such as eBird that compile records that do not originate from standardized programs). All confirmed breeding records were selected. For other types of records, a multiple occupancy criterion, useful to increase confidence that individuals use the habitats in which they were detected, was established at a scale relevant to Eastern Whip-poor-will populations. This was determined to be the 10 x 10 km atlas square because this is consistent with available data (e.g., breeding bird atlas projects, land cover datasets and standardized national grid systems) but also because it is a representative scale to capture the species' needs. It is at this scale that disturbance regimes (both natural and anthropogenic) operate to provide the supply of nesting and foraging habitats necessary to support self-sustaining populations. Consequently, the habitat occupancy criteria for the Eastern Whip-poor-will are met within an atlas square when records from the 2001 breeding season (21st May - 15th August) or thereafter consist of at least :

- One record of confirmed breeding;
OR
- Two records during a single year or from distinct years, where at minimum one record consists of probable breeding;
OR
- Two possible breeding records during a single year in combination with at least one possible breeding record from any other year;
OR
- Five possible breeding records during a single year or from different years.

⁸ Standard breeding evidence definitions used in Canadian breeding bird atlases can be found at www.bsc-eoc.org/norac/atlascodes.htm

⁹ Searching for confirmed breeding activity by this species would not be advisable as it would likely result in inadvertent destruction or disturbance of the nest and eggs

The data used to identify the critical habitat in the present recovery strategy are from 2001 to 2014, inclusively depending on the data available within each region. Since many habitat types occupied by the Eastern Whip-poor-will are dynamic (e.g., a clear cut forest can become too dense for nesting or foraging within 10-15 years under optimal conditions), records older than 2001 will need to be validated to determine if suitable nesting and foraging habitats are still available and if the species still occupies the area.

7.1.2 Habitat Suitability

This criterion refers to the biophysical attributes of habitats in which individuals may carry out breeding (e.g., courtship, territory defense, nesting) and foraging activities in Canada (Table 5). Breeding and foraging habitats form a mosaic. Depending on the habitat attributes available locally, nesting and foraging habitats overlap to various degrees (e.g., individuals forage in wetlands with sufficient perches but do not nest in them, whereas individuals may nest and forage in young regenerating forests). The criteria to determine the extent of suitable habitat vary accordingly. The United States Fish and Wildlife Service (2002) developed a habitat model for the Eastern Whip-poor-will in the northeastern United States. It suggests that when forests with a dense tree cover are used for nesting purposes, only the first 30 m from the edge needs to be included as the species is not associated to the forest interior. On the other side of the spectrum, when habitats can only be used for foraging purposes because the vegetation structure or other characteristics (e.g., drainage) are unsuitable for nesting, data from Rand's (2014) study suggest that it is sufficient to incorporate the habitat up to 1,250 m of the edge with nesting habitats, i.e. the maximum movement observed.

Table 5. Description of the Biophysical Attributes of Suitable Habitats for the Eastern Whip-poor-will.

Components of habitat suitability	Biophysical Attributes
Regional context	Forests (e.g., deciduous, mixedwood, coniferous, treed wetlands) and open habitats (e.g., shrublands, fallow fields, regeneration following fires or clear-cuts, rock and sand outcrops; shrubby wetlands) form a mosaic
Habitats suitable for both nesting and foraging	<ul style="list-style-type: none"> – Forests with sparse to moderate^l tree cover or open habitats <li style="text-align: center;">AND – Sparse to moderate shrub and herbaceous cover <li style="text-align: center;">AND – Well-drained soils (e.g., sand, sandy-loam) <li style="padding-left: 20px;">➤ <i>Within an atlas square, includes all corresponding areas of 3 ha^m or more</i>
Habitats suitable for nesting only <i>[must be adjacent to foraging habitats]</i>	<ul style="list-style-type: none"> – Forests with a dense tree cover <li style="text-align: center;">AND – Sparse to moderate shrub and herbaceous cover <li style="text-align: center;">AND – Well-drained soils (e.g., sand, sandy-loam) <li style="padding-left: 20px;">➤ <i>Within an atlas square, includes all corresponding areas up to 30 m on the interior side of the forest edge</i>
Habitats suitable for foraging only <i>[must be adjacent to nesting habitats]</i>	<ul style="list-style-type: none"> – Forests with sparse tree cover or open habitats <li style="text-align: center;">AND – Dense shrub cover <li style="text-align: center;">AND – Soil drainage is deficient <li style="padding-left: 20px;">➤ <i>Within an atlas square, includes all corresponding areas up to 1,250 m from the edge with suitable nesting habitat</i> <li style="text-align: center;">OR – Agricultural land with scattered shrubs or trees (e.g., hedgerows) that can be used as perches <li style="padding-left: 20px;">➤ <i>Within an atlas square, includes all corresponding areas up to 1,250 m from the edge with suitable nesting habitat</i>

^l Sparse : <25% ; Moderate : 25-75% ; Dense : > 75%

^m Minimum territory size for the Eastern Whip-poor-will (Cink 2002).

7.1.3 Application of the Critical Habitat Criteria

Critical habitat for the Eastern Whip-poor-will is partially identified in this recovery strategy. It corresponds to the areas of suitable nesting and/or foraging habitats within all 10 x 10 km atlas squares meeting the occupancy criteria. Each of these 10 x 10 km

atlas squares is a critical habitat unit. Due to the dynamic nature of some of the habitat components required by the species, no detailed critical habitat mapping is provided. It is also unknown, at this time, to what degree the objective of maintaining the area of occupancy above 3000 km² is met. The identification of critical habitat is considered partial due to limited survey data availability in certain areas of the Eastern Whip-poor-will's distribution and knowledge gaps related to the importance of landscape-level habitat attributes.

Appendix A (Tables A-1 to A-4 and Figures A-1 to A-4) presents the units within which critical habitat is found for the Eastern Whip-poor-will in Canada. Because they both originate from breeding bird atlas squares, critical habitat units for this species are the same as the 10 x 10 km standardized UTM squares used by Environment and Climate Change Canada to indicate the general geographic location of critical habitat (Yellow squares in figures in Appendix A). Overall, 198 critical habitat units are identified for the Eastern Whip-poor-will, including 32 in Manitoba, 110 in Ontario, 51 in Quebec, and five in the Maritimes (all in New Brunswick). More detailed information on the location of critical habitat to support protection of the species and its habitat may be requested, on a need-to-know basis, by contacting Environment and Climate Change Canada's Recovery Planning section at : ec.planificationduretablissement-recoveryplanning.ec@canada.ca.

Any anthropogenic structure (e.g., houses, paved surfaces) and any other areas that do not have the biophysical attributes of suitable habitat are not identified as critical habitat.

7.2 Schedule of Studies to Identify Critical Habitat

Table 6. Schedule of Studies to Identify Critical Habitat

Description of Activity	Rationale	Timeline
Determine habitat characteristics (e.g., spatial scale, configuration) that influence the selection and quality of nesting and foraging habitat for the Eastern Whip-poor-will in Canada	Clarify critical habitat identification criteria and add a landscape component	2018-2023
Conduct standardized nocturnal breeding surveys where: <ul style="list-style-type: none"> - Records since 2001 partially meet the occupancy criteria; - Records date before 2001 and continued occupancy needs to be verified - Insufficient or no surveys have been conducted (e.g., less accessible areas) 	Additional critical habitat identified to meet the short term population and distribution objectives	2018-2028
Identify areas in the southern portion of the breeding distribution in Canada where the species could recolonize habitats following their restoration or management. Atlas blocks that have been abandoned between successive atlas projects may serve as a starting point.	Additional critical habitat identified to meet the long term population and distribution objectives	2018-2038

7.3 Activities Likely to Result in the Destruction of Critical Habitat

Understanding what constitutes destruction of critical habitat is necessary for the protection and management of critical habitat. Destruction is determined on a case by case basis. Destruction would result if part of the critical habitat were degraded, either permanently or temporarily, such that it would not serve its function when needed by the species. Destruction may result from a single or multiple activities at one point in time or from the cumulative effects of one or more activities over time (Government of Canada 2009). Activities described in Table 7 are examples of those likely to cause destruction of critical habitat for the species; however, destructive activities are not necessarily limited to those listed. It should be noted that some activities that would result in the destruction of critical habitat if conducted during the breeding season could also generate the open habitats necessary for foraging as well as nesting habitats in the following years (dynamic habitat mosaic).

Table 7. Examples of Activities Likely to Destroy Critical Habitat for the Eastern Whip-poor-will.

Description of Activity	Description of Effect	Details of Effect
Intensification of agricultural practices (e.g., conversion of extensive cultures to more intensive crops)	Loss or degradation of suitable foraging habitats; Removal of conditions that allow for sufficient prey populations; Fragmentation of habitat; Reduced insect prey populations through the use of pesticides or herbicides on the crops and adjacent habitats	Applicable at all times in critical habitat units (10 x 10 km squares) where the forest cover is already low (e.g., <25%) and if the effect is permanent; Applicable at all times in critical habitat units if activity would result in forest cover falling below 25% and if the effect is permanent; Applicable at all times if biophysical attributes would become unavailable or available in insufficient amounts at the time they are needed by the species.
Conversion of forests to agricultural lands	Loss or degradation of suitable habitats for nesting and/or foraging; Removal of conditions that allow for sufficient prey populations; Fragmentation of habitat	
Energy development and mineral extraction (e.g., pipelines, energy corridors, resource extraction, dams)		
Construction of housing units and other urban infrastructures (e.g., commercial and industrial buildings, playgrounds, roads)		
Forest management (e.g., clearing of shrubs to install tubes in a sugarbush; maintenance in plantations)	Degradation of suitable nesting habitats in dense forests through impacts on the shrub and herbaceous layers required for the nest site and for roosting	
Maintenance of human-made features (e.g., utility rights-of-way, energy corridors military ranges and fields, airports) through vegetation clearing activities	Degradation of suitable nesting habitats through the reduction/elimination of the shrub and herbaceous layers required for nest site placement and loss of perches needed for roosting and foraging; Reduced insect prey populations through the use of herbicides (less insect producing habitats)	These activities, if conducted outside of the breeding season ⁿ , may not be considered as habitat destruction. When possible, these activities could be conducted in successive steps to maintain partial shrub cover.

ⁿ Eastern Whip-poor-will general nesting period in Canada is estimated to extend from mid-May to the end of July (Rousseu and Drolet, unpublished data).

The 25% forest cover (i.e. 2,500 ha) figure within each 10 x 10 km atlas square within which Eastern Whip-poor-will records occur was selected as a value under which activities, if they were to occur, are more likely to result in the abandonment of the remaining suitable habitats. The following considerations were taken into account :

- The Eastern Whip-poor-will is generally considered a bird of forested landscapes;
- Each individual may require nesting and foraging resources within home ranges as large as 500 ha (Rand 2014). Many of the atlas squares meeting the occupancy criteria had 3 or more individuals;
- Individuals often occupy transitional habitats (e.g., a partial cut forest can become too dense for nesting or foraging within 10-15 years under optimal conditions) making it necessary to maintain alternative nesting and foraging sites over areas larger than those occupied in any given year. This consideration is particularly important in atlas squares where forest cover is already under 25% and alternative breeding habitats may no longer be available ;
- Regions with higher forest cover are more likely to sustain Eastern Whip-poor-will populations over the long term (more suitable habitats, reduced impact of some threats);
- An analysis shows that 10 x 10 km atlas squares meeting the occupancy criteria described in section 7.1.1 have a mean forest cover of 55% in Ontario (range = 0.4-98%); 42% in Quebec (range = 8-78%) and 74% in the Maritimes (range = 62-91%). This implies that there is some flexibility to maintain habitat supply above 25% within most 10 x 10 km squares within which critical habitat is found.

The 25% forest cover figure should be viewed as a general indication of the level of additional activities that could take place within each 10 x 10 km square rather than a true threshold. Other considerations, including the configuration of remaining forest habitats (both within each 10 x 10 km and any neighbouring atlas square where the habitat is contiguous) and the level of fragmentation are also key elements. Some site-level biophysical attributes (e.g., well-drained soils such as sand deposits) may also be available in limited amounts.

8. Measuring Progress

The performance indicators presented below provide a way to define and measure progress toward achieving the population and distribution objectives.

- In the short term (2018-2028), declining population trends have been halted or reversed to a point where the Canadian population of the Eastern Whip-poor-will have declined no more than 10% and the areas of occupied habitats are maintained at 3000 km² or more.
- In the long term (after 2028), a positive 10-year trend, measured by BBS and other available data (e.g., targeted surveys), is achieved (i.e., the population is increasing) and the species has started gradually recolonizing former areas in the southern portion of the Canadian range.

9. Statement on Action Plans

One or more action plans for the Eastern Whip-poor-will will be posted on the Species at risk public registry before the end of 2023.

10. References

- Aide, T. M., Clark, M. L., Grau, H. R., López-Carr, D., Levy, M. A., Redo, D., Bonilla-Moheno, M., Riner, G., Andrade-Núñez, M. J. and Muñiz, M. 2013. Deforestation and reforestation of latin America and the Caribbean (2001–2010). *Biotropica* 45: 262–271.
- Allombert, S., S. Stockton, and J.-L. Martin. 2005. A natural experiment on the impact of overabundant deer on forest invertebrates. *Conservation Biology* 19(6):1917-1929.
- Anodon, J.D., O. E. Sala and F.T. Maestre. 2014. Climate change will increase savannas at the expense of forests and treeless vegetation in tropical and subtropical Americas. *Journal of Ecology* (102) : 1363-1373.
- Atlas des oiseaux nicheurs du Québec. 2014. Données obtenues de la part des bureaux de l'Atlas des oiseaux nicheurs du Québec. Regroupement QuébecOiseaux, Service canadien de la faune d'Environnement Canada et Études d'Oiseaux Canada. Québec, Québec, Canada. Online : www.atlas-oiseaux.qc.ca
- Bélanger, L., M. Grenier and S. Deslandes, 1999. Bilan des habitats et de l'occupation du sol dans la vallée du Saint-Laurent. Environnement Canada, Service canadien de la faune, Région du Québec.
- Benton, T. G., D. M. Bryant, L. Cole and H. Q. P. Crick. 2002. Linking agricultural practice to insect and bird populations: a historical study over three decades. *Journal of Applied Ecology* 39: 673-687.
- Blancher, P., M. D. Cadman, B. A. Pond, A. R. Couturier, E. H. Dunn, C. M. Francis and R. S. Rempel. 2007. Changes in bird distributions between atlases. In M. D. Cadman, D. A. Sutherland, G. G. Beck, D. Lepage and A. R. Couturier (eds.). *Atlas of the breeding birds of Ontario, 2001-2005*. Pp. 32-48. Bird Studies Canada, Environment Canada, Ontario Field Ornithologists, Ontario Ministry of Natural Resources, and Ontario Nature, Toronto (Ontario).
- Blancher, P.J., R.D. Phoenix, D.S. Badzinski, M.D. Cadman, T.L. Crewe, C.M. Downes, D. Fillman, C.M. Francis, J. Hughes, D.J.T. Hussell, D. Lepage, J.D. McCracken, D.K. McNicol, B.A. Pond, R.K. Ross, R. Russell, L.A. Venier and R.C. Weeber. 2009. Population trend status of Ontario's forest birds. *The Forestry Chronicle* 85(2):184-201.

Bock, C.E. and Z.F. Jones. 2004. Avian habitat evaluation: should counting birds count? *Frontiers in Ecology and Environment* 2: 403–410.

Boettner, G. H., J.S. Elkinton and C. J. Boettner. 2000. Effects of a biological control introduction on three nontarget native species of nonnative Saturniid moths. *Conservation Biology* 14(6): 1798-1806.

Both, C., C. A. Van Turnhout, R. G. Bijlsma, H. Siepel, A. J. Van Strien and R. P. Foppen. 2009. Avian population consequences of climate change are most severe for long-distance migrants in seasonal habitats. *Proceedings of the Royal Society B: Biological Sciences* 277: 1259-1266.

Both, C., S. Bouwhuis, C. Lessells and M.E. Visser. 2006. Climate change and population declines in a long-distance migratory bird. *Nature* 441(7089): 81-83.

Broeckart, M. and S. Bédard. 2012. Rapport sur la validation du modèle de l'habitat des engoulevants au Québec, la précision de l'habitat convenable pour l'Engoulevant bois-pourri et la détermination de routes d'inventaires permanentes. Regroupement QuébecOiseaux. 9 p.

Broeckart, M. 2012. Réalisation cartographique de la potentialité de l'habitat de l'Engoulevant bois-pourri au Québec. Regroupement QuébecOiseaux. 115 pp.

Brooks, D. R., J. E. Bajer, S. J. Clark, D. T. Monteith, C. Andrews, S. J. Corbett, D. A. Beaumont and J. W. Chapman. 2012. Large carabid beetle declines in a United Kingdom monitoring network increases evidence for a widespread loss in insect biodiversity. *Journal of Applied Ecology* 49(5): 1009-1019.

Brown, C. R. and M. B. Brown. 2000. Weather-mediated natural selection on arrival time in Cliff Swallows (*Petrochelidon pyrrhonota*). *Behavioral Ecology and Sociobiology* 47(5): 339-345.

Bushman, E.S. and G.D. Therres. 1988. Habitat management guidelines for forest interior breeding birds of coastal Maryland. Maryland Department of Natural Resources. Wildlife Technical Publication 88-1. 50 p.

Cadman, M. D., D.A. Sutherland, G. G. Beck, D. LePage and A.R. Couturier, eds. 2007. Atlas of the breeding birds of Ontario, 2001-2005. Bird Studies Canada, Environment Canada, Ontario Field Ornithologists, Ontario Ministry of Natural Resources, and Ontario Nature, Toronto, xxii + 706 p.

Cadman, M. D., Sutherland, D. A. and F. M. Helleiner (eds). 1987. Atlas of the Breeding Birds of Ontario. University of Waterloo Press. Waterloo, Ontario. 617 p.

Chesser, R. T., Banks, R.C., Barker, F. K., Cicero, C., Dunn, J. L., Kratter, A. W., Lovette, I. J., Rasmussen, P. C., Remsen, J. V., Jr, Rising, J. D., Stotz, D. F. and K. Winker. 2010. Fifty-first supplement to the American ornithologists' Union *Check-list of North American Birds*. The Auk. 127(3):726-744.

Chesser, R. T., R.C. Banks, F. K. Barker, C. Cicero, J. L. Dunn, A. W. Kratter, I. J. Lovette, P. C. Rasmussen, J. V. Remsen, J. D. Jr Rising, D. F. Stotz, and K. Winker. 2012. Fifty-third supplement to the American ornithologists' Union *Check-list of North American Birds*. The Auk. 129(3):573-588.

Chiron, F., R. Chargé, R. Julliard, F. Jiguet and A. Muratet. 2014. Pesticide doses, landscape structure and their relative effects on farmland birds. *Agriculture, Ecosystems and Environment* 185:153–160.

Cink, C. L. 2002. Eastern Whip-poor-will (*Antrostomus vociferus*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America. Online: <http://bna.birds.cornell.edu/bna/species/620> (Accessed September, 2014).

Colburn, T., F.S. Vom Saal and A.M. Soto. 1993. Developmental effects of endocrine-disrupting chemicals in wildlife and humans. *Environmental Health Perspectives* 101: 378-384.

Cooper, R.J. 1981. Relative abundance of Georgia Caprimulgids based on call-counts. *Wilson Bulletin* 93: 363-371.

Cornell Lab of Ornithology. 2007. Eastern Whip-poor-will. Online: http://www.allaboutbirds.org/guide/eastern_whip-poor-will/lifehistory (Accessed April, 2013).

COSEWIC. 2009. COSEWIC assessment and status report on the Whip-poor-will *Caprimulgus vociferus* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vi + 28 pp.

Côté, S. D., T. P. Rooney, J.-P. Tremblay, C. Dussault and D. Waller. 2004. Ecological impacts of deer overabundance. *Annual Review of Ecology, Evolution, and Systematics* 3: 113-147.

Cunningham, H.M, K. Chaney, R.B. Bradbury and A. Wilcox. 2004. Non-inversion tillage and farmland birds: a review with special reference to the UK and Europe. *Ibis* 146(2): 192–202.

Cyr, A. and J. Larivée. 1995. Atlas saisonnier des oiseaux du Québec. Société du Loisir Ornithologique. 714 p.

DeGraaf, R. M., W. M. Healy and R. T. Brooks. 1991. Effects of thinning and deer browsing on breeding birds in New England oak woodlands. *Forest Ecology and Management* 41:179-191.

de Groot, W. J., M. D. Flannigan and A. S. Cantin. 2013. Climate change impacts on future boreal fire regimes. *Forest Ecology and Management* 294: 35-44.

Dendroica Environnement et Faune. 2013. Inventaire de huit espèces d'oiseaux en péril en Outaouais en vue de désigner leur habitat essentiel : 1ere année du suivi. Présenté au Service canadien de la faune Environnement Canada. 23 p.

Dionne, M., C. Maurice, J. Gauthier and F. Shaffer. 2008. Impact of hurricane Wilma on migrating birds: The case of the Chimney Swift. *The Wilson Journal of Ornithology* 120(4): 784-792.

Dirzo, R., H. S. Young, M. Galetti, G. Ceballos, N. J. Isaac and B. Collen. 2014. Defaunation in the Anthropocene. *Science* 345(6195): 401-406.

Eastman, J. 1991. Whip-poor-will, pp. 252-253 in Brewer, R., G.A. McPeck and R.J. Adams Jr., eds. *The Atlas of Breeding Birds of Michigan*. East Lansing, MI: Michigan State University Press, East Lansing, Michigan. 594pp.

eBird Canada. 2014. Bird Observations on eBird website Version 1.0. Audubon and Cornell Lab of Ornithology. Online: <http://ebird.org/ebird/canada/> (Accessed April, 2014).

English, P. and M.A. Conboy. 2013. Developing stewardship prescriptions for Whip-poor-wills (*Antrostomus vociferous*) in eastern Ontario: SARSF Final Report. Ontario Ministry of Natural Resources. Peterborough, Ontario. 14 p.

Environment Canada. 2011. Presence and levels of priority pesticides in selected Canadian aquatic ecosystems. Environment Canada, Water Science and Technology Directorate. Gatineau, Québec.

Environment Canada. 2014a. North American Breeding Bird Survey – Canadian Results and Analysis Website version 3.00. Environment Canada, Gatineau, Quebec. Online : <http://ec.gc.ca/reom-mbs/default.asp?lang=En&n=0D74F35F-1> (Accessed June, 2014).

Environment Canada. 2014b. Avoidance Guidelines related to Incidental Take of Migratory Birds in Canada. Online: <http://www.ec.gc.ca/paom-itmb>

Erskine, A.J. 1992. *Atlas of Breeding Birds of the Maritime Provinces*. Nimbus Publishing Ltd. The Nova Scotia Museum. 270 p.

Flannigan, M., B. Stocks, M. Turetsky and M. Wotton. 2009. Impacts of climate change on fire activity and fire management in the circumboreal forest. *Global Change Biology* 15(3): 549-560.

Foster, G. N. 1991. Conserving insects of aquatic and wetland habitats, with special reference to beetles. Pages 237-262 *In* The conservation of insects and their habitats. 15th Symposium of the Royal Entomological Society of London. Academic Press. London, UK.

Gallizioli, S. 1979. Effects of livestock grazing on wildlife. Presented at the 10th Annual joint Meeting of the western Section, the Wildlife Society and the California-Nevada Chapter, American Fisheries Society. Online: <http://www.rangebiome.org/cowfree/gallizioli/EffectsOfLivestockGrazing79.htm> (Accessed November, 2014).

Garlapow, R. M. 2007. Whip-poor-will prey availability and foraging habitat: implications for management in pitch pine / scrub oak barrens habitats. University of Massachusetts - Amherst. Masters Theses. Paper 27. 58 p.

Gauthier, J. and Y. Aubry (eds). 1996. The breeding birds of Quebec: Atlas of the breeding birds of southern Quebec. Province of Quebec Society for the Protection of Birds and Canadian Wildlife Service, Montreal. xviii + 1,302 pp.

Gill, R. 2000. The Impact of deer on woodland biodiversity. Forestry Commission Information Note 36. Forestry Commission, Edinburgh.

Girardin, M. P., A. A. Ali, C. Carcaillet, O. Blarquez, C. Hély, A. Terrier, A. Genries and Y. Bergeron. 2013. Vegetation limits the impact of a warm climate on boreal wildfires. *New Phytologist* 199(4): 1001-1011.

Godfrey, W.E. 1986. Birds of Canada. National Museum of Natural Sciences, Ottawa, 595 pp.

Goetsch, C., J. Wigg, A. A. Royo, T. Ristau, and W. P. Carson. 2011. Chronic over browsing and biodiversity collapse in a forest understory in Pennsylvania: Results from a 60 year-old deer exclusion plot. *Journal of the Torrey Botanical Society* 138 (2):220-224.

Goldstein, M. I., T. E. J. Lacher, B. Woodbridge, M. J. Bechard, S. B. Canavelli, M. E. Saccagnini, G. P. Cobb, E. J. Scollon, R. Tribolet and M. J. Hooper. 1999. Monocrotophos-induced mass mortality of Swainson's Hawks in Argentina, 1995-96. *Ecotoxicology* 8:201-214.

Gornish, E. S. and J. M. Tylianakis. 2013. Community shifts under climate change: Mechanisms at multiple scales. *American Journal of Botany* 100(7):1422-1434.

Goulson, D. 2013. Review: An overview of the environmental risks posed by neonicotinoid insecticides. *Journal of Applied Ecology* 50(4):977-987.

Goulson, D. 2014. Pesticides linked to bird declines. *Nature* 511: 295-296.

Government of Canada. 2009. *Species at Risk Act Policies, Overarching framework [Draft]. Species at Risk Act Policy and Guideline Series*. Environment Canada. Ottawa. 38 pp.

Graveland, J. 1998. Effects of acid rain on bird populations. *Environmental Review* 6:41-54.

Hallmann, C. A., R. P. Foppen, C. A. van Turnhout, H. de Kroon and E. Jongejans. 2014. Declines in insectivorous birds are associated with high neonicotinoid concentrations. *Nature* 511: 341-343.

Hansen, M., P. Potapov, R. Moore, M. Hancher, S. Turubanova, A. Tyukavina, D. Thau, S. Stehman, S. Goetz and T. Loveland. 2013. High-resolution global maps of 21st-century forest cover change. *Science* 342(6160): 850-853.

Hawley, D. M., K. K. Hallinger and D. A. Cristol. 2009. Compromised immune competence in free-living Tree Swallows exposed to mercury. *Ecotoxicology* 18(5): 499-503.

Hobson, K. A., A. G. Wilson, S. L. Van Wilgenburg and E. M. Bayne. 2013. An estimation of nest loss in Canada due to industrial forestry operations. *Avian Conservation and Ecology* 8(2):5.

Huber, D. G. and J. Gullede. 2011. *Extreme weather and climate change: understanding the link, managing the risk*. Center for Climate and Energy Solutions, Arlington, VA.

Hunt, P. D. 2010. Whip-poor-will territory mapping at two New Hampshire sites. Audubon Society of New Hampshire, Concord, NH, USA. 9 p.

Jobin, B., C. Latendresse, A. Baril, C. Maisonneuve and D. Côté. 2014. A half-century analysis of landscape dynamics in southern Québec, Canada. *Environmental Monitoring and Assessment* 186: 2215–2229.

Jones, G. A., K. E. Sieving and S. K. Jacobson. 2005. Avian diversity and functional insectivory on North-Central Florida Farmlands. *Proceedings: Conservation Biology* 19(4):1234-1245.

Jones, T. and W. Cresswell. 2010. The phenology mismatch hypothesis: are declines of migrant birds linked to uneven global climate change? *Journal of Animal Ecology* 79(1): 98-108.

Jorgenson, T. and A. Foster. 2007. *Grazing Forested Rangeland*. Government of Saskatchewan. Online: <http://www.agriculture.gov.sk.ca/Default.aspx?DN=4aabe84e-ea79-417a-ace2-e629ffe95cb3> (Accessed December, 2013).

- Keller, R. H., L. Xie, D. B. Buchwalter, K. E. Franzreb and T. R. Simons. 2014. Mercury bioaccumulation in Southern Appalachian birds, assessed through feather concentrations. *Ecotoxicology* 23(2): 304-316.
- Klemens, J. A., R. G. Harper, J. A. Frick, A. P. Capparella, H. B. Richardson and M. J. Coffey. 2000. Pattern of organochlorine pesticide contamination in neotropical migrant passerines in relation to diet and winter habitat. *Chemosphere* 41:1107-1113.
- Larivée, J. 2013. Étude des populations d'oiseaux du Québec (Version 2013-04-11) [base de données]. Rimouski, Québec : Regroupement QuébecOiseaux.
- Laughlin, S. B. and D. P. Kibbe. 1985. The atlas of breeding birds of Vermont. University Press of New England, Hanover, New Hampshire, USA.
- Leonard, H. J. 1987. Natural resources and economic development in Central America: a regional environmental profile. For the International Institute for Environment and Development.
- Lewis, T. and G. C. Dibley. 1970. Air movement near windbreaks and a hypothesis of the mechanism of the accumulation of airborne insects. *Annals of Applied Biology* 66:477-484.
- Lewis, T. and J. W. Stephenson. 1966. The permeability of artificial windbreaks and the distribution of flying insects in the leeward sheltered zone. *Annals of Applied Biology* 58:355-363.
- Longcore, T. and C. Rich. 2004. Ecological light pollution. *Frontiers in Ecology and the Environment* 2(4):191-198.
- Machtans, C. S., C. H. Wedeles et E. M. Bayne. 2013. A first estimate for Canada of the number of birds killed by colliding with building windows. *Avian Conservation and Ecology* 8(2): 6.
- Main, A. R., J. V. Headley, K. M. Peru, N. L. Michel, A. J. Cessna and C. A. Morrissey. 2014. Widespread use and frequent detection of neonicotinoid insecticides in wetlands of Canada's prairie pothole region. *PLoS one* 9(3): e92821.
- Maritime Breeding Bird Atlas. 2013. Bird Studies Canada and Environment Canada – CWS – Atlantic region. Online: <http://www.mba-aom.ca/>
- Masek, J. G., W. B. Cohen, D. Leckie, M. A. Wulder, R. Vargas, B. de Jong, S. Healey, B. Law, R. Birdsey and R. Houghton. 2011. Recent rates of forest harvest and conversion in North America. *Journal of Geophysical Research: Biogeosciences* (2005–2012) 116(G4).

Medler, M. D. 2008. Whip-poor-will *Caprimulgus vociferus*. Pages 310-311 dans McGowan, K. J. and K. Corwin (eds). The second atlas of breeding birds in New York State. Cornell University Press. Ithaca, New York. 687 p.

Mehlman, D.W., S.E. Mabey, D.N. Ewert, C. Duncan, B. Abel, D. Cimprich, R.D. Sutter, and M. Woodrey. 2005. Conserving stopover sites for forest-dwelling migratory landbirds. *Auk* 122(4):1281-1290.

Mills, A. M. 1986. The influence of moonlight on the behaviour of goatsuckers (*Caprimulgidae*). *The Auk* 103(2):370-378.

Mills, A. M. 2007. Whip-poor-will, pp. 312-313 in Cadman, M.D., D.A. Sutherland, G.G. Beck, D. LePage and A.R. Couturier, eds. Atlas of the breeding birds of Ontario, 2001-2005. Bird Studies Canada, Environment Canada, Ontario Field Ornithologists, Ontario Ministry of Natural Resources, and Ontario Nature, Toronto, xxii + 706 pp.

Mineau, P. and C. Palmer. 2013. The impact of the nation's most widely used insecticides on birds: neonicotinoid insecticides and birds. American Bird Conservancy, Washington, DC.

Møller, A. P., D. Rubolini and E. Lehikoinen. 2008. Populations of migratory bird species that did not show a phenological response to climate change are declining. *Proceedings of the National Academy of Sciences* 105(42): 16195-16200.

NatureServe. 2014. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Online: <http://www.natureserve.org/explorer> (Accessed March, 2014).

Nebel, S., A. Mills, J. D. McCracken and P. D. Taylor. 2010. Declines of aerial insectivores in North America follow a geographic gradient. *Avian Conservation and Ecology – Écologie et conservation des oiseaux* 5(2): 1.

NFD. 2014. National Forestry Database. Natural Resources Canada and Canadian Forest Service, Ottawa, ON. Online: http://nfdp.ccfm.org/index_e.php. (Accessed August 20, 2014).

Nocera, J. J., J. M. Blais, D. V. Beresford, L. K. Finity, C. Grooms, L. E. Kimpe, K. Kyser, N. Michelutti, M. W. Reudink and J. P. Smol. 2012. Historical pesticide applications coincided with an altered diet of aerially foraging insectivorous chimney swifts. *Proceedings: Biological Sciences* 279(1740):3114-3120.

North American Bird Conservation Initiative US Committee. 2010. The State of the Birds 2010 - Report on Climate Change, United States of America. US Department of the Interior, Washington, D.C.

Ontario Ministry of Natural Resources. 2009. Whip-poor-will (*Caprimulgus vociferus*). Queen's Printer for Ontario. 2 p.

Osborne, C. E., D. C. Evers, M. Duron, N. Schoch, D. Yates, D. Buck, O. P. Lane, and J. Franklin. 2011. Mercury Contamination within Terrestrial Ecosystems in New England and Mid-Atlantic States: Profiles of Soil, Invertebrates, Songbirds, and Bats. Report BRI 2011-09. Submitted to The Nature Conservancy – Eastern New York Chapter. Biodiversity Research Institute, Gorham, Maine. 100 p.

Palmer-Ball, B.L. Jr. 1996. The Kentucky breeding bird atlas. Univ. Press of Kentucky, Lexington.

Paquette, S.R., F. Pelletier, D. Garant and M. Bélisle. 2014. Severe recent decrease of adult body mass in a declining insectivorous bird population. *Proceedings of the Royal Society B* 281: 9 p.

Parish, T., K.H. Lakhani, and T.H. Sparks. 1995. Modelling the relationship between bird population variables and hedgerow, and other field margin attributes. II. Abundance of individual species and of groups of similar species. *Journal of Applied ecology* 32(2):362-371.

Partners in Flight Science Committee. 2013. Population Estimates Database. Online: <http://rmbo.org/pifpopestimates> (Accessed October, 2013).

Patric, J. H. and J. D. Helvey. 1986. Some effects of grazing on soil and water in the Eastern forest. United States Department of Agriculture, Northeastern Forest Experiment Station. Broomall, Pennsylvania. 11 p.

Peck, G.K. and R.D. James. 1983. Breeding birds of Ontario, nidiology and distribution, Vol. 1: nonpasserines. Royal Ontario Museum, Toronto, 321 pp.

Peterson, W. R. and W. R. Meservey. 2003. Massachusetts breeding bird atlas. Massachusetts Audubon Society, Lincoln, Massachusetts, USA.

Pollard, E. and A.S. Cooke. 1994. Impact of muntjac deer *Muntiacus reevesi* on egg-laying sites of the white admiral butterfly *Ladoga camilla* in a cambridgeshire wood. *Biological Conservation* 70(2):189-191.

Price, P. W., R. F. Denno, M. D. Eubanks, D. L. Finke and I. Kaplan. 2011. Insect ecology: behavior, populations and communities. Cambridge University Press. New York, NY. 812 pp.

Radeloff, V. C., R. B. Hammer. and S. I. Stewart. 2005. Rural and sub-urban sprawl in the U.S. Midwest from 1940 to 2000 and its relation to forest fragmentation. *Conservation Biology* 19:793-805.

Ramankutty, N., E. Heller and J. Rhemtulla. 2010. Prevailing myths about agricultural abandonment and forest regrowth in the United States. *Ann. Assoc. Am. Geogr.* 100:502-512.

Rand, G.J. 2014. Home range use, habitat selection, and stress physiology of Eastern Whip-poor-wills (*Antrostomus vociferus*) at the northern edge of their range. Msc. Thesis, Trent University. Peterborough, Ontario. 66 pp.

Reese, J.G. 1996. Whip-poor-will, pp. 194-195 in C.S. Robbins, ed. *Atlas of the Breeding Birds of Maryland and the District of Columbia*. Univ. of Pittsburgh Press, Pittsburgh, PA.

Robbins, C.S. 1994. Whip-poor-will. In C.R. Foss (ed.) *Atlas of breeding birds in New Hampshire*. Audubon Society of New Hampshire, Concord, New Hampshire. 414 p.

Robinson, L., J. P. Newell and J. M. Marzluff. 2005. Twenty-five years of sprawl in the Seattle region: Grown management responses and implications for conservation. *Landscape and Urban Planning* 71:51-72.

Robinson, R. A. and W. J. Sutherland. 2002. Post-war changes in arable farming and biodiversity in Great Britain. *Journal of Applied Ecology* 39:157-176.

Rooney, T. P. 2009. High white-tailed deer densities benefit graminoids and contribute to biotic homogenization of forest ground-layer vegetation. *Plant ecology*, 202(1):103-111.

Rousseu, F. and B. Drolet. Unpublished data. The nesting phenology of birds in Canada. Technical Report Series. Canadian Wildlife Service, Environment Canada.

Roy, L. and M. Bombardier. 1996. Whip-poor-will. Pages 626-629 in Gauthier, J. and Y. Aubry (eds). *The breeding birds of Quebec: Atlas of the breeding birds of Southern Quebec*. Province of Quebec Society for the Protection of Birds and Canadian Wildlife Service, Montreal. xviii + 1,302 p.

Russell, F.L., D.B. Zippin and/ N.L. Fowler. 2001. Effects of white-tailed deer (*Odocoileus virginianus*) on plants, plant populations and communities: a review. *American Midl/ Naturalist* 146:1-26.

Sager, T. A. 1997. Organochlorine pesticide contamination in new world passerines. Honors Project, Paper 10. Illinois Wesleyan University.

Saino, N., R. Ambrosini, D. Rubolini, J. von Hardenberg, A. Provenzale, K. Hüppop, O. Hüppop, A. Lehikoinen, E. Lehikoinen and K. Rainio. 2011. Climate warming, ecological mismatch at arrival and population decline in migratory birds. *Proceedings of the Royal Society B: Biological Sciences* 278(1707): 835-842.

Sandilands, A. P. 2010. Birds of Ontario: habitat requirements, limiting factors and status. Vol. II, Nonpasserines: Shorebirds through Woodpeckers. University of British Columbia Press. 387 pp.

Santner, S. 1992. Whip-poor-will. Pages 172-173. In Atlas of breeding birds in Pennsylvania. Braunong, D. W., (Ed.) University of Pittsburgh, PA.

Sauer, J. R., G. W. Pendleton and B. G. Peterjohn. 1996. Evaluating causes of population change in North American insectivorous songbirds. *Conservation Biology* 10(2):465-478.

Scheuhammer, A. M., M. W. Meyer, M. B. Sandheinrich and M. W. Murray. 2007. Effects of environmental methylmercury on the health of wild birds, mammals, and fish. *AMBIO: A Journal of the Human Environment* 36(1): 12-19.

Smith, A.R. 1996. Atlas of Saskatchewan birds. Canadian Wildlife Service, Natural History Society, Regina, 456 pp.

Strode, P. K. 2003. Implications of climate change for North American wood warblers (Parulidae). *Global Change Biology* 9(8): 1137-1144.

Sun, H., W. Forsythe and/ N. Waters. 2007. Modeling urban land use change and urban sprawl: Calgary, Alberta, Canada. *Networks and Spatial Economics* 7:353-376.

Taper, M. L., K. Böhning-Gaese and J. H. Brown. 1995. Individualistic responses of bird species to environmental change. *Oecologia* 101(4):478-486.

Tozer, D.C., J.C. Hoare, J.E. Inglis, J. Yaraskavitch, H. Kitching and S. Dobbyn. 2014. Clearcut with seed trees in red pine forests associated with increased occupancy by Eastern Whip-poor-wills. *Forest Ecology and Management* 330: 1-7.

Turpak, J.M., D.T. Jones-Farrand, F.R. Thompson III, D.J. Twedt and W.B. Uihlein III. 2009. Multiscale habitat suitability index models for priority landbirds in the Central Hardwoods and West Gulf Coastal Plain/Ouachitas Bird Conservation regions. United States Department of Agriculture - Forest Service. General Technical Report NRS-49. Online: <http://www.nrs.fs.fed.us/mwg-internal/de5fs23hu73ds/progress?id=vDvypziygX>

Tyler, W. M. 1940. Eastern Whip-poor-will. Pages 163-183 in Life histories of North American cuckoos, goatsuckers, hummingbirds and their allies. (Bent, A. C., Ed.) U.S. Natl. Mus. Bull. 176p.

U.S. Bureau of Land Management. 1978. Grass creek: oil and gas leasing environmental assessment record. U.S. Bureau of Land Management, Worland District, WY.

United States Fish and Wildlife Service. 2002. Whip-poor-will habitat model.
http://www.fws.gov/r5gomp/gom/habitatstudy/metadata2/whip-poor-will_model.htm

Van Wilgenburg, S.L., K.A. Hobson, E.M. Bayne and N. Koper. 2013. Estimated avian nest loss associated with oil and gas exploration and extraction in the Western Canadian Sedimentary Basin. *Avian Conservation and Ecology* 8(2): 9.

Verboom, B. and K. Spoelstra. 1999. Effects of food abundance and wind on the use of tree lines by an insectivorous bat, *Pipistrellus pipistrellus*. *Canadian Journal of Zoology*, 77:1393-1401.

Visser, M.E., L. J. M. Holleman and P. Gienapp. 2006. Shifts in caterpillar biomass phenology due to climate change and its impact on the breeding biology of an insectivorous bird. *Oecologia*. 147(1):164-172.

Wagner, D. L. 2012. Moth decline in the Northeastern United States. *News of the Lepidopterists' Society* 54(2): 52-56.

Wickramasinghe, L. P., S. Harris, G. Jones and N. V. Jennings. 2004. Abundance and species richness of nocturnal insects on organic and conventional farms: effects of agricultural intensification on bat foraging. *Conservation Biology* 18 (5):1283-1292.

Wilson, M.D. 2003. Distribution, abundance, and home range of the Whip-poor-will (*Caprimulgus vociferus*) in a managed forest landscape. MS. Thesis. College of William and Mary, Williamsburg, VA.

Wilson, M.D. and B.D. Watts. 2008. Landscape configuration effects on distribution and abundance of Whip-poor-wills. *The Wilson Journal of Ornithology* 120(4):778-783.

Wilson, S.G. 1985. Summer distribution of Whip-poor-wills in Minnesota. *Loon* 57:6-8.

Xing, Z., L. Chow, H. Rees, F. Meng, S. Li, B. Ernst, G. Benoy, T. Zha and L. M. Hewitt. 2013. Influences of sampling methodologies on pesticide-residue detection in stream water. *Archives of Environmental Contamination and Toxicology* 64(2): 208-218.

Appendix A: Critical Habitat for the Eastern Whip-poor-will in Canada

Table A-1. Critical habitat units (10 x 10 km standardized UTM squares) within which critical habitat for the Eastern Whip-poor-will is found in Manitoba. Critical habitat occurs where the criteria described in Section 7.1 are met.

10 x 10 km Standardized Grid Square ID ^o	UTM Grid Square Coordinates ^p		Land Tenure ^q
	Easting	Northing	
14ULE48	340000	5980000	Other federal Lands and Non-federal Lands
14UMC93	490000	5730000	Non-federal Lands
14UMD66	460000	5860000	Non-federal Lands
14UMD76	470000	5860000	Non-federal Lands
14UMD87	480000	5870000	Non-federal Lands
14UMD95	490000	5850000	Non-federal Lands
14UMF80	480000	6000000	Non-federal Lands
14UNB56	550000	5660000	Non-federal Lands
14UNB87	580000	5670000	Other federal Lands and Non-federal Lands
14UNB96	590000	5660000	Non-federal Lands
14UNC23	520000	5730000	Non-federal Lands
14UNC53	550000	5730000	Non-federal Lands
14UND16	510000	5860000	Non-federal Lands
14UPA66	660000	5560000	Non-federal Lands
14UPA76	670000	5560000	Non-federal Lands
14UPA85	680000	5550000	Non-federal Lands
14UPB70	670000	5600000	Non-federal Lands
14UPB79	670000	5690000	Non-federal Lands

10 x 10 km Standardized Grid Square ID ^o	UTM Grid Square Coordinates ^p		Land Tenure ^q
	Easting	Northing	
14UPB97	690000	5670000	Other federal Lands and Non-federal Lands
14UPV64	660000	5440000	Non-federal Lands
14UPV89	680000	5490000	Non-federal Lands
14UQA11	710000	5510000	Non-federal Lands
15UTQ94	290000	5440000	Non-federal Lands
15UTS96	289996	5660000	Non-federal Lands
15UUR08	300000	5580000	Non-federal Lands
15UUR18	310000	5580000	Non-federal Lands
15UUR29	320000	5590000	Non-federal Lands
15UUS05	300000	5650000	Non-federal Lands
15UUS06	300000	5660000	Non-federal Lands
15UUS21	320000	5610000	Non-federal Lands
15UUS35	330000	5650000	Non-federal Lands
15UUT37	330000	5770000	Non-federal Lands
Total: 32 critical habitat units			

^o Based on the standard UTM Military Grid Reference System (see <http://www.nrcan.gc.ca/earth-sciences/geography/topographic-information/maps/9789>), where the first 2 digits represent the UTM Zone, the following letter represents the UTM Row, the next 2 letters indicate the 100 km x 100 km Standardized UTM grid, followed by 2 digits to represent the 10 x 10 km Standardized UTM grid containing all or a portion of the critical habitat unit. This unique alphanumeric code is based on the methodology produced from the Breeding Bird Atlases of Canada (See <http://www.bsc-eoc.org/> for more information on breeding bird atlases).

^p The listed coordinates are a cartographic representation of where critical habitat can be found, presented as the southwest corner of the 10 x 10 km Standardized UTM grid squares containing all or a portion of the critical habitat unit. The coordinates may not fall within critical habitat and are provided as a general location only.

^q Land tenure is provided as an approximation of the types of land ownership that exist at the critical habitat units and should be used for guidance purposes only. Accurate land tenure will require cross referencing critical habitat boundaries with surveyed land parcel information.

Table A-2. Critical habitat units (10 x 10 km standardized UTM squares) within which critical habitat for the Eastern Whip-poor-will is found in Ontario. Critical habitat occurs where the criteria described in Section 7.1 are met.

10 x 10 km standardized Grid Square ID ^r	UTM Grid Square Coordinates ^s		Land tenure ^t
	Easting	Northing	
15UVQ11	410000	5410000	Non-federal Lands
15UVQ21	420000	5410000	Non-federal Lands
15UVQ31	430000	5410000	Non-federal Lands
16UCV80	380000	5400000	Other Federal Land and Non-federal Lands
16TGS25	720000	5150000	Other Federal Land and Non-federal Lands
17TLM66	360000	5160000	Non-federal Lands
17TMG28	420000	4680000	Non-federal Lands
17TMG38	430000	4680000	Non-federal Lands
17TMH28	420000	4780000	Other Federal Land and Non-federal Lands
17TMH38	430000	4780000	Non-federal Lands
17TMK61	460000	4910000	Non-federal Lands
17TMK69	460000	4990000	Other Federal Land and Non-federal Lands
17TMK78	470000	4980000	Non-federal Lands
17TMK79	470000	4990000	Non-federal Lands
17TML40	440000	5000000	Other Federal Land and Non-federal Lands
17TML50	450000	5000000	Other Federal Land and Non-federal Lands
17TML60	460000	5000000	Other Federal Land and Non-federal Lands
17TML69	460000	5090000	Other Federal Land and Non-federal Lands
17TML70	470000	5000000	Other Federal Land and Non-federal Lands
17TML71	470000	5010000	Other Federal Land and Non-federal Lands

10 x 10 km standardized Grid Square ID ^r	UTM Grid Square Coordinates ^s		Land tenure ^t
	Easting	Northing	
17TMM04	400000	5140000	Non-federal Lands
17TNH32	530000	4720000	Non-federal Lands
17TNH41	540000	4710000	Federal Protected Area (Big Creek National Wildlife Area) and Non-federal Lands
17TNH42	540000	4720000	Non-federal Lands
17TNH52	550000	4720000	Non-federal Lands
17TNH61	560000	4710000	Federal Protected Area (Long Point National Wildlife Area) and Non-federal Lands
17TNH69	560000	4790000	Non-federal Lands
17TNH71	570000	4710000	Federal Protected Area (Long Point National Wildlife Area) and Non-federal Lands
17TNH79	570000	4790000	Non-federal Lands
17TNK88	580000	4980000	Non-federal Lands
17TNK96	590000	4960000	Federal Protected Area (Georgian Bay Islands National Park) and Non-federal Lands
17TNL37	530000	5070000	Other Federal Land and Non-federal Lands
17TNL43	540000	5030000	Non-federal Lands
17TNL62	560000	5020000	Non-federal Lands
17TNL70	570000	5000000	Non-federal Lands
17TNL87	580000	5070000	Non-federal Lands
17TNL88	580000	5080000	Non-federal Lands
17TNN86	580000	5260000	Non-federal Lands
17TPH35	630000	4750000	Non-federal Lands
17TPJ77	670000	4870000	Non-federal Lands

10 x 10 km standardized Grid Square ID ^r	UTM Grid Square Coordinates ^s		Land tenure ^t
	Easting	Northing	
17TPJ87	680000	4870000	Non-federal Lands
17TPJ98	690000	4880000	Non-federal Lands
17TPK17	610000	4970000	Other Federal Land and Non-federal Lands
17TPK25	620000	4950000	Other Federal Land and Non-federal Lands
17TPK46	640000	4960000	Non-federal Lands
17TPK53	650000	4930000	Other Federal Land and Non-federal Lands
17TPK54	650000	4940000	Non-federal Lands
17TPK55	650000	4950000	Non-federal Lands
17TPK56	650000	4960000	Non-federal Lands
17TPK57	650000	4970000	Non-federal Lands
17TPK64	660000	4940000	Other Federal Land and Non-federal Lands
17TQJ38	730000	4880000	Non-federal Lands
17TQK03	700000	4930000	Other Federal Land and Non-federal Lands
17TQK14	710000	4940000	Non-federal Lands
17TQK23	720000	4930000	Other Federal Land and Non-federal Lands
17TQK24	720000	4940000	Non-federal Lands
17TQK26	720000	4960000	Non-federal Lands
17TQK32	730000	4920000	Non-federal Lands
17TQK34	730000	4940000	Other Federal Land and Non-federal Lands
17TQL07	700000	5070000	Non-federal Lands
18TTP77	270000	4780000	Non-federal Lands
18TTQ63	261440	4930000	Non-federal Lands
18TTQ64	261808	4940000	Non-federal Lands

10 x 10 km standardized Grid Square ID ^r	UTM Grid Square Coordinates ^s		Land tenure ^t
	Easting	Northing	
18TTQ74	270000	4940000	Other Federal Land and Non-federal Lands
18TTQ75	270000	4950000	Non-federal Lands
18TTQ93	290000	4930000	Non-federal Lands
18TTR74	270000	5040000	Non-federal Lands
18TTR88	280000	5080000	Non-federal Lands
18TTR96	290000	5060000	Other Federal Land and Non-federal Lands
18TTR98	290000	5080000	Other Federal Land and Non-federal Lands
18TUP25	320000	485000	Other Federal Land and Non-federal Lands
18TUP46	340000	486000	Federal Protected Area (Prince Edward Point National Wildlife Area) and Non-federal Lands
18TUP88	380000	488000	Non-federal Lands
18TUQ03	300000	4930000	Non-federal Lands
18TUQ28	320000	4980000	Non-federal Lands
18TUQ43	340000	4930000	Non-federal Lands
18TUQ54	350000	4940000	Non-federal Lands
18TUQ63	360000	4930000	Non-federal Lands
18TUQ67	360000	4970000	Non-federal Lands
18TUQ72	370000	4920000	Non-federal Lands
18TUQ73	370000	4930000	Non-federal Lands
18TUQ74	370000	4940000	Other Federal Land and Non-federal Lands
18TUQ75	370000	4950000	Non-federal Lands
18TUQ80	380000	4900000	Other Federal Land and Non-federal Lands
18TUQ82	380000	4920000	Other Federal Land and Non-federal Lands

10 x 10 km standardized Grid Square ID ^r	UTM Grid Square Coordinates ^s		Land tenure ^t
	Easting	Northing	
18TUQ83	380000	4930000	Other Federal Land and Non-federal Lands
18TUQ92	390000	4920000	Other Federal Land and Non-federal Lands
18TUQ93	390000	4930000	Other Federal Land and Non-federal Lands
18TUQ95	390000	4950000	Other Federal Land and Non-federal Lands
18TUR04	300000	5040000	Non-federal Lands
18TUR06	300000	5060000	Non-federal Lands
18TUR16	310000	5060000	Non-federal Lands
18TUR17	310000	5070000	Non-federal Lands
18TUR18	310000	5080000	Other Federal Land and Non-federal Lands
18TUR25	320000	5050000	Non-federal Lands
18TUR28	320000	5080000	Other Federal Land and Non-federal Lands
18TUS00	300000	5100000	Other Federal Land and Non-federal Lands
18TVQ01	400000	4910000	Non-federal Lands
18TVQ03	400000	4930000	Other Federal Land and Non-federal Lands
18TVQ04	400000	4940000	Other Federal Land and Non-federal Lands
18TVQ05	400000	4950000	Other Federal Land and Non-federal Lands
18TVQ12	410000	4920000	Non-federal Lands
18TVQ22	420000	4920000	Other Federal Land and Non-federal Lands
18TVQ27	420000	4970000	Federal Protected Area (Rideau Migratory Bird Sanctuary), Other Federal Land and Non-federal Lands
18TVQ37	430000	4970000	Federal Protected Area (Rideau Migratory Bird Sanctuary), Other Federal Land and Non-federal Lands
18TVQ38	430000	4980000	Other Federal Land and Non-federal Lands

10 x 10 km standardized Grid Square ID ^r	UTM Grid Square Coordinates ^s		Land tenure ^t
	Easting	Northing	
18TVR10	410000	5000000	Non-federal Lands
18TVR12	410000	5020000	Non-federal Lands
18TVR22	420000	5020000	Other Federal Land and Non-federal Lands
18TVR82	480000	5020000	Non-federal Lands
Total: 110 critical habitat units			

^r Based on the standard UTM Military Grid Reference System (see <http://www.nrcan.gc.ca/earth-sciences/geography/topographic-information/maps/9789>), where the first 2 digits represent the UTM Zone, the following letter represents the UTM Row, the next 2 letters indicate the 100 km x 100 km Standardized UTM grid, followed by 2 digits to represent the 10 x 10 km Standardized UTM grid containing all or a portion of the critical habitat unit. This unique alphanumeric code is based on the methodology produced from the Breeding Bird Atlases of Canada (See <http://www.bsc-eoc.org/> for more information on breeding bird atlases).

^s The listed coordinates are a cartographic representation of where critical habitat can be found, presented as the southwest corner of the 10 x 10 km Standardized UTM grid squares containing all or a portion of the critical habitat unit. The coordinates may not fall within critical habitat and are provided as a general location only.

^t Land tenure is provided as an approximation of the types of land ownership that exist at the critical habitat units and should be used for guidance purposes only. Accurate land tenure will require cross referencing critical habitat boundaries with surveyed land parcel information.

Table A-3. Critical habitat units (10 x 10 km standardized UTM squares) within which critical habitat for the Eastern Whip-poor-will is found in Quebec. Critical habitat occurs where the criteria described in Section 7.1 are met.

10 x 10 km Standardized Grid Square ID ^u	UTM Grid Square Coordinates ^v		Land Tenure ^w
	Easting	Northing	
17TPN26	620000	5260000	Non-federal Lands
17UPP45	640000	5350000	Non-federal Lands
17UPP55	650000	5350000	Non-federal Lands
18TUR38	330000	5070000	Other Federal Lands and Non-federal Lands
18TUR47	340000	5070000	Other Federal Lands and Non-federal Lands
18TUR85	380000	5050000	Non-federal Lands
18TUR93	390000	5030000	Other Federal Lands and Non-federal Lands
18TUR95	390000	5050000	Other Federal Lands and Non-federal Lands
18TUS33	330000	5130000	Other Federal Lands
18TVR19	410000	5040000	Other Federal Lands and Non-federal Lands
18TVR14	410000	5090000	Other Federal Lands and Non-federal Lands
18TVR24	420000	5040000	Other Federal Lands and Non-federal Lands
18TVR26	420000	5060000	Non-federal Lands
18TVR28	420000	5080000	Non-federal Lands
18TVR35	430000	5050000	Non-federal Lands
18TVR55	450000	5050000	Non-federal Lands
18TVT10	410000	5200000	Non-federal Lands
18TWQ59	550000	4990000	Non-federal Lands
18TWQ88	580000	4980000	Other Federal Lands and Non-federal Lands
18TWQ89	580000	4990000	Non-federal Lands
18TWQ98	590000	4980000	Other Federal Lands and Non-federal Lands

10 x 10 km Standardized Grid Square ID ^u	UTM Grid Square Coordinates ^v		Land Tenure ^w
	Easting	Northing	
18TWR56	550000	5060000	Non-federal Lands
18TWR66	560000	5060000	Non-federal Lands
18TWR90	590000	5000000	Non-federal Lands
18TWR98	590000	5080000	Non-federal Lands
18TXQ09	600000	4990000	Non-federal Lands
18TXR00	600000	5000000	Other Federal Lands and Non-federal Lands
18TXR10	610000	5000000	Non-federal Lands
18TXR29	620000	5090000	Other Federal Lands and Non-federal Lands
18TXR46	640000	5060000	Non-federal Lands
18TXR50	650000	5000000	Non-federal Lands
18TXR51	650000	5010000	Other Federal Lands and Non-federal Lands
18TXR52	650000	5020000	Other Federal Lands and Non-federal Lands
18TXR62	660000	5020000	Non-federal Lands
18TXR73	670000	5030000	Non-federal Lands
18TXR89	680000	5090000	Non-federal Lands
18TXR96	690000	5060000	Non-federal Lands
18TXS74	670000	5140000	Other Federal Lands and Non-federal Lands
18UXV60	660000	5400000	Other Federal Lands and Non-federal Lands
18TYR08	700000	5080000	Non-federal Lands
18TYR18	710000	5080000	Non-federal Lands
18TYR19	710000	5090000	Non-federal Lands
18TYR28	720000	5080000	Non-federal Lands
18TYS10	710000	5100000	Other Federal Lands and Non-federal Lands

10 x 10 km Standardized Grid Square ID ^u	UTM Grid Square Coordinates ^v		Land Tenure ^w
	Easting	Northing	
18TYS21	720000	5110000	Non-federal Lands
18TYS31	730000	5110000	Non-federal Lands
19TBM71	270000	5110000	Non-federal Lands
19TBM72	270000	5120000	Non-federal Lands
19TCM06	300000	5160000	Non-federal Lands
19TCM26	320000	5160000	Non-federal Lands
19UDP43	440000	5330000	Other Federal Lands and Non-federal Lands
Total: 51 critical habitat units			

^u Based on the standard UTM Military Grid Reference System (see <http://www.nrcan.gc.ca/earth-sciences/geography/topographic-information/maps/9789>), where the first 2 digits represent the UTM Zone, the following letter represents the UTM Row, the next 2 letters indicate the 100 km x 100 km Standardized UTM grid, followed by 2 digits to represent the 10 x 10 km Standardized UTM grid containing all or a portion of the critical habitat unit. This unique alphanumeric code is based on the methodology produced from the Breeding Bird Atlases of Canada (See <http://www.bsc-eoc.org/> for more information on breeding bird atlases).

^v The listed coordinates are a cartographic representation of where critical habitat can be found, presented as the southwest corner of the 10 x 10 km Standardized UTM grid squares containing all or a portion of the critical habitat unit. The coordinates may not fall within critical habitat and are provided as a general location only.

^w Land tenure is provided as an approximation of the types of land ownership that exist at the critical habitat units and should be used for guidance purposes only. Accurate land tenure will require cross referencing critical habitat boundaries with surveyed land parcel information.

Table A-4. Critical habitat units (10 x 10 km standardized UTM squares) within which critical habitat for the Eastern Whip-poor-will is found in New Brunswick. Critical habitat occurs where the criteria described in Section 7.1 are met.

10 x 10 km Standardized Grid Square ID ^x	UTM Grid Square Coordinates ^y		Land Tenure ^z
	Easting	Northing	
19TFL96	690000	5060000	Other Federal Lands and Non-federal Lands
19TGL17	710000	5070000	Other Federal Lands and
19TGL19	710000	5090000	Non-federal Lands
19TFN24	620000	5240000	Non-federal Lands
20TLT01	300000	5210000	Other Federal Lands and Non-federal Lands

^x Based on the standard UTM Military Grid Reference System (see <http://www.nrcan.gc.ca/earth-sciences/geography/topographic-information/maps/9789>), where the first 2 digits represent the UTM Zone, the following letter represents the UTM Row, the next 2 letters indicate the 100 km x 100 km Standardized UTM grid, followed by 2 digits to represent the 10 x 10 km Standardized UTM grid containing all or a portion of the critical habitat unit. This unique alphanumeric code is based on the methodology produced from the Breeding Bird Atlases of Canada (See <http://www.bsc-eoc.org/> for more information on breeding bird atlases).

^y The listed coordinates are a cartographic representation of where critical habitat can be found, presented as the southwest corner of the 10 x 10 km Standardized UTM grid squares containing all or a portion of the critical habitat unit. The coordinates may not fall within critical habitat and are provided as a general location only.

^z Land tenure is provided as an approximation of the types of land ownership that exist at the critical habitat units and should be used for guidance purposes only. Accurate land tenure will require cross referencing critical habitat boundaries with surveyed land parcel information.




Figure A-1. Overview of maps A-1.1 to A-1.11 representing the critical habitat for the Eastern Whip-poor-will in Manitoba. Critical habitat for the Eastern Whip-poor-will in Manitoba occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-1.1. Critical habitat for the Eastern Whip-poor-will in Manitoba occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-1.2. Critical habitat for the Eastern Whip-poor-will in Manitoba occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-1.3. Critical habitat for the Eastern Whip-poor-will in Manitoba occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-1.4. Critical habitat for the Eastern Whip-poor-will in Manitoba occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-1.5. Critical habitat for the Eastern Whip-poor-will in Manitoba occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-1.6. Critical habitat for the Eastern Whip-poor-will in Manitoba occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-1.7. Critical habitat for the Eastern Whip-poor-will in Manitoba occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-1.8. Critical habitat for the Eastern Whip-poor-will in Manitoba occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-1.9. Critical habitat for the Eastern Whip-poor-will in Manitoba occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-1.10. Critical habitat for the Eastern Whip-poor-will in Manitoba occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-1.11. Critical habitat for the Eastern Whip-poor-will in Manitoba occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2. Overview of maps A-2.1 to A-2.23 representing the critical habitat for the Eastern Whip-poor-will in Ontario. Critical habitat occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.1. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.2. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.3. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.4. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.5. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.6. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.7. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.8. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.9. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.10. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.11. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.12. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.13. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.14. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.15. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.16. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.17. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.18. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.19. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.20. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.21. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.22. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-2.23. Critical habitat for the Eastern Whip-poor-will in Ontario occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-3. Overview of maps A-3.1 to A-3.12 representing the critical habitat for the Eastern Whip-poor-will in Quebec. Critical habitat for the Eastern Whip-poor-will in Quebec occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-3.1. Critical habitat for the Eastern Whip-poor-will in Quebec occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-3.2. Critical habitat for the Eastern Whip-poor-will in Quebec occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-3.3. Critical habitat for the Eastern Whip-poor-will in Quebec occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-3.4. Critical habitat for the Eastern Whip-poor-will in Quebec occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-3.5. Critical habitat for the Eastern Whip-poor-will in Quebec occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-3.6. Critical habitat for the Eastern Whip-poor-will in Quebec occurs within the 10 x 10 km Standardized UTM grid squares indicated (squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-3.7. Critical habitat for the Eastern Whip-poor-will in Quebec occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-3.8. Critical habitat for the Eastern Whip-poor-will in Quebec occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-3.9 Critical habitat for the Eastern Whip-poor-will in Quebec occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-3.11. Critical habitat for the Eastern Whip-poor-will in Quebec occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-3.12. Critical habitat for the Eastern Whip-poor-will in Quebec occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-4. Overview of maps A-4.1 to A-4.3 representing the critical habitat for the Eastern Whip-poor-will in New Brunswick. Critical habitat for the Eastern Whip-poor-will in New Brunswick occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-4.1. Critical habitat for the Eastern Whip-poor-will in New Brunswick occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-4.2. Critical habitat for the Eastern Whip-poor-will in New Brunswick occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.




Figure A-4.3. Critical habitat for the Eastern Whip-poor-will in New Brunswick occurs within the 10 x 10 km Standardized UTM grid squares indicated (yellow squares), where the criteria and methodology set out in Section 7.1 are met. This standardized national grid system indicates the general location within which critical habitat is found.

Appendix B. Effects on the Environment and Other Species

A strategic environmental assessment (SEA) is conducted on all SARA recovery planning documents, in accordance with the [Cabinet Directive on the Environmental Assessment of Policy, Plan and Program Proposals](#)¹⁰. The purpose of a SEA is to incorporate environmental considerations into the development of public policies, plans, and program proposals to support environmentally sound decision-making and to evaluate whether the outcomes of a recovery planning document could affect any component of the environment or any of the [Federal Sustainable Development Strategy](#)'s¹¹ (FSDS) goals and targets.

Recovery planning is intended to benefit species at risk and biodiversity in general. However, it is recognized that strategies may also inadvertently lead to environmental effects beyond the intended benefits. The planning process based on national guidelines directly incorporates consideration of all environmental effects, with a particular focus on possible impacts upon non-target species or habitats. The results of the SEA are incorporated directly into the strategy itself, but are also summarized below in this statement.

All species that depend on aerial insects for prey such as bats, swallows, and specifically, bird species at risk including: the Chimney Swift (*Chaetura pelagica*), the Common Nighthawk (*Cordeiles minor*), the Olive-sided Flycatcher (*Contopus cooperi*), the Acadian Flycatcher (*Empidonax virescens*) may benefit from the recommended approaches for Eastern Whip-poor-will, namely by increasing the availability of insect producing habitats. These species are also important for integrated pest management.

Nonetheless, some species, including other species at risk, may prefer different forest conditions than the Eastern Whip-poor-will, which needs open areas for nesting. Recovery actions for the species must be integrated with best practices for other songbird species, especially where the best practices may conflict.

The possibility that the present recovery strategy inadvertently generates negative effects on the environment and on other species was considered. The majority of recommended actions are non-intrusive in nature, including surveys and outreach. We conclude that the present recovery strategy is unlikely to produce significant negative effects.

¹⁰ www.ceaa.gc.ca/default.asp?lang=En&n=B3186435-1

¹¹ www.ec.gc.ca/dd-sd/default.asp?lang=En&n=CD30F295-1